

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/122347/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Wan, Zhenzhu, Algeo, Thomas J., Gensel, Patricia G., Scheckler, Stephen E., Stein, William E., Cressler, Walter L., Berry, Christopher M. ORCID: <https://orcid.org/0000-0001-9521-5618>, Xu, Honghe, Rowe, Harold D. and Sauer, Peter E. 2019. Environmental influences on the stable carbon isotopic composition of Devonian and Early Carboniferous land plants. *Palaeogeography, Palaeoclimatology, Palaeoecology* 531 (A) , 109100. 10.1016/j.palaeo.2019.02.025 file

Publishers page: <http://dx.doi.org/10.1016/j.palaeo.2019.02.025>
<<http://dx.doi.org/10.1016/j.palaeo.2019.02.025>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Environmental influences on the stable carbon isotopic composition of Devonian and Early Carboniferous land plants

Zhenzhu Wan^a, Thomas J. Algeo^{a,b,c,*}, Patricia G. Gensel^d, Stephen E. Scheckler^e, William E. Stein^f, Walter L. Cressler III^g, Christopher M. Berry^h, Honghe Xuⁱ, Harold D. Rowe^j, and Peter E. Sauer^k

^a *Department of Geology, University of Cincinnati, Cincinnati, OH 45221-0013, USA*

^b *State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan 430074, China*

^c *State Key Laboratory of Geological Processes and Mineral Resources, China University of Geosciences, Wuhan 430074, China*

^d *Department of Biology, University of North Carolina, Chapel Hill, NC 27599 USA*

^e *Depts. of Biological Sciences and Geosciences, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0406, USA*

^f *Department of Biological Sciences, SUNY Binghamton, Binghamton, NY 13901, USA*

^g *Francis Harvey Green Library, West Chester University, West Chester, PA 19383, USA*

^h *Department of Earth Sciences, Cardiff University, Cardiff CF10 3YE, Wales, UK*

ⁱ *State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Nanjing 210008, China*

^j *Bureau of Economic Geology, University of Texas at Austin, Austin, TX 78713-8924 USA*

^k Department of Earth and Atmospheric Sciences, Indiana University, Bloomington, IN 47405-2208, USA

*Corresponding author: thomas.algeo@uc.edu

Abstract

Systematic analysis of the stable carbon isotopic composition of fossil land plants ($\delta^{13}\text{C}_\text{p}$) has the potential to offer new insights regarding paleoclimate variation and plant-environment interactions in early terrestrial ecosystems. $\delta^{13}\text{C}_\text{p}$ was measured for 190 fossil plant specimens belonging to 10 genera of Early to Late Devonian age (*Archaeopteris*, *Drepanophycus*, *Haskinsia*, *Leclercqia*, *Pertica*, *Psilophyton*, *Rhacophyton*, *Sawdonia*, *Tetraxylopteris*, and *Wattieza*) and 2 genera of Early Carboniferous age (*Genselia* and *Rhodeopteridium*) collected from sites located mainly in the Appalachian Basin (22-30 °S paleolatitude). For the full carbon-isotopic dataset ($n = 309$), $\delta^{13}\text{C}_\text{p}$ ranges from -20.3 ‰ to -30.5 ‰ with a mean of -25.5 ‰, similar to values for modern C_3 land plants. In addition to a secular trend, $\delta^{13}\text{C}_\text{p}$ exhibits both intra- and intergeneric variation. Intrageneric variation is expressed as a small (mean 0.45 ‰) ^{13}C -enrichment of leaves and spines relative to stems that may reflect differential compound-specific compositions. Intergeneric variation is expressed as a much larger (to ~5 ‰) spread in the mean $\delta^{13}\text{C}_\text{p}$ values of coeval plant genera that was probably controlled by taxon-specific habitat preferences and local environmental humidity. Among Early Devonian taxa, *Sawdonia* yielded the most ^{13}C -depleted values (-27.1 ± 1.7 ‰), reflecting lower water-use efficiency that was probably related to growth in wetter habitats, and *Leclercqia*, *Haskinsia*, and *Psilophyton* yielded the most ^{13}C -

enriched values (-23.0 ± 1.6 ‰, -22.3 ± 1.3 ‰, and -24.8 ± 1.6 ‰, respectively), reflecting higher water-use efficiency probably related to growth in drier habitats.

Keywords: water-use efficiency; early plants; canopy effect; *Archaeopteris*; *Drepanophycus*; *Sawdonia*

1. Introduction

The Devonian was a period of revolutionary changes in terrestrial floras, landscapes, and continental climates (Algeo et al., 1995, 2001; Gibling and Davies, 2012; Gibling et al., 2014).

Although bryophytes (non-vascular plants) had appeared during the mid-Ordovician and tracheophytes (vascular plants) by the late Silurian (Wellman et al., 2003; Steemans et al., 2009), the continents remained sparsely vegetated until the Devonian. That period witnessed major morphological adaptations and diversifications among vascular land plants as well as order-of-magnitude increases in the complexity and geographic extent of terrestrial ecosystems (Gensel and Edwards, 2001; Taylor et al., 2009). Early Devonian land plants were mostly small (heights of no more than a few tens of centimeters), shallowly rooted where known, and limited to moist lowland habitats (Xue et al., 2016). By the end-Devonian, vascular land plants had evolved leaves, complex root systems, secondary supporting tissues (leading to tree stature), and desiccation-resistant reproductive propagules (seeds) that facilitated their spread into drier upland habitats (Mosbrugger, 1990; Driese et al., 1997; Gerrienne et al., 2004; Meyer-Berthaud et al., 2010; Hao and Xue, 2013a; Silvestro et al., 2015). [Note: “upland” is a general term signifying land lying above the level of local stream flow and flooding.] These changes

66 resulted in an increase in organic carbon burial, a rise of O₂ and a drawdown of CO₂ in the
67 atmosphere, and global climatic cooling, setting the stage for the Late Paleozoic Ice Age that
68 began in the Early Carboniferous (Algeo et al., 1995, 2001; Berner, 1997; Le Hir et al., 2011).

69 The stable carbon isotopic composition of plants ($\delta^{13}\text{C}_p$) differs from that of their carbon
70 source (atmospheric CO₂) owing to photosynthetic fractionation (Farquhar et al., 1982, 1989).
71 Modern C₃ plants fix carbon via the Calvin-Benson cycle with an average fractionation of ca.
72 -20 ‰, yielding organic material with a $\delta^{13}\text{C}$ value of -26 ± 5 ‰ (Raven et al., 2004). Devonian
73 vascular land plants are thought to have been exclusively C₃ plants (Taylor et al., 2009), and
74 their carbon isotopic compositions are therefore likely to have been controlled by the same
75 climatic and environmental factors as for modern C₃ plants (e.g., temperature, humidity, and
76 atmospheric $p\text{CO}_2$ and $\delta^{13}\text{C}_{\text{CO}_2}$). To date, the carbon isotopic compositions of specific Devonian
77 terrestrial plant fossils (as opposed to bulk sedimentary organic carbon) have been investigated
78 in only a few studies of fungi (Boyce et al., 2007), lichens (Fletcher et al., 2004), liverworts
79 (Graham et al., 2010), and vascular plants (Beerling et al., 2002; Feng et al., 2014).

80 The present study is the first systematic investigation of carbon isotopic variation among
81 a large group of Devonian and Early Carboniferous vascular land plants and represents by far
82 the largest such analytical dataset to date. We made 309 measurements of $\delta^{13}\text{C}_p$ (p = plant) on
83 190 unique specimens of fossil plants belonging to 12 widespread genera of Early Devonian to
84 Early Carboniferous age (Table S1). Our goals in this study were to (1) characterize $\delta^{13}\text{C}_p$
85 variation for each fossil plant taxon, (2) investigate time-independent $\delta^{13}\text{C}_p$ variation within and
86 among taxa that might reflect large-scale geographic (e.g., paleolatitudinal) or local
87 environmental influences, and (3) use these patterns of variation to draw inferences concerning

the ecology of individual early land plant taxa. Here, we show that there are significant differences in $\delta^{13}\text{C}_\text{p}$ between the 12 genera of this study that are probably of primary (*in vivo*) origin, and that were most likely related to local humidity and habitat preferences among early vascular land plants.

2. Fossil plant taxonomy and paleoecology

2.1. Fossil plant taxa

The fossil plants analyzed in this study belong to 12 different genera of tracheophytes, representing early members of the lycophyte and euphyllophyte lineages (Fig. 1A). All of the fossil plants in this study have been the subject of at least partial morphological reconstructions (Fig. 1B). Although many fossil specimens were identified to the species level, we conducted our evaluation of $\delta^{13}\text{C}_\text{p}$ variation at the genus level owing to (1) lack of species assignments for about half of the individual specimens, (2) uncertainty in some species-level assignments, and (3) the impracticability of analyzing carbon-isotopic patterns in a dataset with excessive splitting into species. Among the 12 genera analyzed, 10 have overlapping ranges extending through part or all of the Devonian (*Archaeopteris*, *Drepanophycus*, *Haskinsia*, *Leclercqia*, *Pertica*, *Psilophyton*, *Rhacophyton*, *Sawdonia*, *Tetraxylopteris*, and *Wattieza*) and two are entirely of Early Carboniferous age (*Genselia* and *Rhodeopteridium*; Fig. 2). A brief summary of the key anatomical features and paleoecologic/geographic characteristics of each genus follows:

Sawdonia (Dawson) Hueber is one of the early zosterophylloids, belonging to the Lycophyta. This plant is pseudomonopodially branched with circinate tips; reniform sporangia are borne along the sides of some stems and anatomy consists of an exarch haplostele (Hueber,

1971; Rayner, 1983; Gensel et al., 1975; Taylor et al., 2009; Gensel and Berry, 2016). It was widespread during the Early Devonian (Gensel, 1992; Kenrick and Crane, 1997; Gensel and Berry, 2016) but is also found in the Late Devonian (Hueber and Grierson, 1961). Our dataset includes both *S. ornata* and *S. acanthotheca*, although most specimens were not distinguished at the species level.

Drepanophycus Göppert belongs to the Drepanophycaceae, sister to other lineages of Lycopsidea (Kenrick and Crane, 1997). Stems range from several millimeters to centimeters in diameter and from several centimeters to a meter in length. Leaves are unbranched falcate microphylls up to several millimeters long with a single prominent middle vascular thread, and they were arranged spirally or randomly on the stem. Reniform sporangia occur on short stalks between the leaves (Li and Edwards, 1995; Li et al., 2000). However, this genus exhibits considerable variation in leaf morphology and may be in need of revision. This taxon had a creeping to erect and sometimes dichotomizing rhizomatous growth habit. It existed through most of the Devonian Period and is found in eastern Canada, northeastern USA, China, Russia, and Europe (Stubblefield and Banks, 1978; Gensel and Berry, 2001; Xu et al., 2013). About half of the specimens used in the present study belong to *D. spinaeformis*, and the remainder were not distinguished at the species level.

Leclercqia Banks, Bonamo and Grierson is a member of the Protolepidodendraceae (Lycopphyta) (Gensel and Kasper, 2005; Gensel and Albright, 2006). It is a slender, herbaceous plant, with distinctive leaves that typically exhibit five divisions and that bear sporangia on their upper surface. At least some species exhibit hook-shaped leaves that may have allowed a vine-like climbing habit (Xu et al., 2011). *Leclercqia* is distinguished in being a homosporous plant

with a ligule; ligules otherwise are found only in heterosporous taxa (Grierson and Bonamo, 1979; Bonamo et al., 1988; Taylor et al., 2009; Benca et al., 2014). It existed from the late Early to the Middle Devonian and has a particularly wide distribution, being found in North America, Europe, Africa, Australia, and China (Banks et al., 1972; Bonamo et al., 1988; Meyer-Berthaud et al., 2003; Taylor et al., 2009; Xu et al., 2011; Prestianni et al., 2012; Benca et al., 2014). It was present mainly in warm, dry climates (Berry, 1994; Meyer-Berthaud et al., 2003), although some studies have claimed a wider environmental range (Xu and Wang, 2008). Our dataset includes both *L. complexa* and *L. andrewsii*, although half of the specimens were not distinguished at the species level.

Haskinsia Grierson and Banks is a herbaceous lycopsid (Bonamo et al., 1988; Xu et al., 2008; Yang et al., 2008; Taylor et al., 2009). It was once regarded as a species of *Drepanophycus* but was later placed with the Protolepidodendrales because of the presence of petiolate deltoid-shaped sporophylls (Grierson and Banks, 1983; Berry and Edwards, 1996; Xu and Berry, 2008). This taxon was widespread during the Middle to early Late Devonian age, being known from North America, South America, Russia, and China; it had wide environmental tolerances, thriving in both warm tropical and cool temperate conditions (Xu et al., 2008). The relatively small number of specimens in our dataset includes *H. hastata*, *H. sagittata*, and specimens not identified at the species level.

Psilophyton (Dawson) Hueber is one of the best-known members of the trimerophyte grade of basal euphyllophytes (Andrews et al., 1968; Banks et al., 1975; Doran et al., 1978; Trant and Gensel, 1985; Stein, 1993; Gerrienne 1995, 1997). The plant consists of a main axis and multiply divided lateral branches, some of which terminate in pairs of fusiform sporangia.

Stem anatomy, where known, is a centrarch haplostele. Intrageneric variation in sporangial length and in the presence and type of emergences is significant (Doran, 1980; Gerrienne, 1995; Taylor et al., 2009). *Psilophyton* was widespread in the Early to earliest Middle Devonian and has been found in the northeastern United States (Maine and New York), Canada (Quebec and New Brunswick), Belgium, the Czech Republic, and Yunnan, China (Taylor et al., 2009; Gensel, 2017). The most common species in our dataset is *P. princeps*, although specimens of *P. crenulatum*, *P. dawsonii*, and *P. forbesii* are also present, as well as some specimens not identified at the species level.

Pertica Kasper and Andrews is also a basal euphyllophyte (Kasper and Andrews, 1972; Granoff et al., 1976). It exhibits a dominant main stem with regularly arranged lateral branches, organized either in a tight spiral or a four ranked/decussate pattern. Lateral branch morphology varies from equally dichotomous to a more pseudomonopodial pattern. These plants bore numerous fusiform sporangia in dense clusters on ultimate branchlets (Kasper and Andrews, 1972; Granoff et al., 1976; Taylor et al., 2009). *Pertica* existed from the Early to the Middle Devonian, with occurrences known from northern Maine, New Brunswick, and Quebec. Our dataset consists mainly of the species *P. varia* along with some specimens not identified at the species level.

Wattieza refers to compression fossils belonging to the order Pseudosporochnales within the class Cladoxylopsida (Leclercq and Banks, 1962; Gensel et al., 2001; Berry and Fairon-Demaret, 2002). Pseudosporochnaleans are all similar in their adult forms, consisting of small to large trees with a main trunk bearing a dense crown of frond-like branches. Instead of leaves, the branches bear densely ramified ultimate appendages with either terminally erect

(*Pseudosporochnus*) or reflexed (*Wattieza*) sporangia. The specimens analyzed for this study, which came from the West Cave Mountain and Steenberg/South Mountain quarries in New York, possessed key characteristics of Pseudosporochnaleans including distinctive clusters of sclerotic cells (Leclercq and Banks, 1962; Gensel et al., 2001; Berry and Fairon-Demaret, 2002) but lacked sporangia. However, a recent study of new South Mountain specimens identified them as *Wattieza* and linked them to the fossil tree trunks known as *Eospermatopteris* (Stein et al., 2007). Pseudosporochnaleans existed from the Middle to early Late Devonian in Europe, Venezuela, North America, and possibly China (Gensel and Andrews, 1984; Taylor et al., 2009).

Rhacophyton Crépin belongs to the order of Rhacophytales *sensu* Taylor et al. (2009; n.b., assigned to Rhacophytaceae or Protopteridales by others). According to Andrews and Phillips (1968), this plant grew to ~1-m-tall bushes, and its foliage consisted of a main axis bearing pairs of lateral branches that fork twice basally. Two of the lateral branches were multiply divided and terminated in elongate sporangia, and the other two extended further and bore higher-order branches. In some cases, the fertile branches were produced as second-order units on first-order laterals. The stem anatomy of the lateral branches consisted of clepsydroid primary xylem surrounded in some instances by a limited amount of secondary xylem. *Rhacophyton* existed during the Late Devonian and has been found in North America, Europe, possibly Bear Island (New Brunswick), and western Siberia (Andrews and Phillips, 1968; Cornet et al., 1977; Dittrich et al., 1983; Cressler, 1999, 2006; Taylor et al., 2009). Our dataset includes a few specimens identified as *R. ceratangium* although the majority were not identified at the species level.

Rhodeopteridium (Presl) Zimmermann (formerly *Rhodea* Presl) is a genus based mainly on foliage compression/impression fossils that is thought to represent an early seed plant (Jennings, 1976). Pinnately compound leaves terminate in barely laminate ends (Read, 1955). Similar fossil leaves with well-preserved petioles are known from early seed ferns of Late Mississippian (Chesterian) age in Illinois (Jennings, 1976). Most specimens are of Early Carboniferous (Tournaisian-Namurian A) age, but it may extend into the basal late Carboniferous. None of the small number of *Rhodeopteridium* specimens in our dataset was identified at the species level.

Tetraxylopteris Beck is an aneurophytalean progymnosperm (Beck, 1957; Bonamo and Banks, 1967; Scheckler and Banks, 1971; Hammond and Berry, 2005). It is characterized by a main stem with extensive secondary xylem and bearing several orders of laterals in a decussate pattern, each with a four-armed vascular strand. In fertile regions, second-order laterals dichotomize twice, and each division bears pinnately arranged clusters of sporangia. *Tetraxylopteris* existed during the Givetian and early Frasnian. To date, it is known from only two areas: the Catskill Delta in New York, and the Campo Chico Formation in northwestern Venezuela. A few of our specimens are assigned to *T. schmidtii* but most were not identified at the species level.

Archaeopteris (Dawson) Stur belongs to the Archaeopteridalean progymnosperms. It was a tall tree similar to some modern conifers, but it had pseudomonopodial branching in the lateral branch system (Beck, 1962) and laminate leaves with helices or decussate patterns of leaf attachment (Scheckler, 1978). Some ultimate clusters of leaves appear cone-like, bearing sporangia on their adaxial surfaces, but all species are thought to have been free-sporing and

heterosporous. *Archaeopteris* is usually preserved as impression and compression fossils, and its fossilized trunks, named *Callixylon* when found separately, exhibit extensive secondary xylem. It ranged from the latest Middle to Late Devonian, with a number of different species being known from localities in North America, Russia, Europe, Morocco, China, Australia, and Colombia (Beck and Wight, 1988; Cressler, 1999, 2006; Meyer-Berthaud et al., 1999; Berry et al., 2000). By the middle of the late Frasnian, monospecific archaeopterid forests had become the dominant vegetation type in lowland areas and coastal settings over a vast geographic area (Algeo et al., 2001). About one third of the 51 specimens in our dataset are assigned to various species, including *A. fissilis*, *A. halliana*, *A. jacksonii*, *A. macilenta*, and *A. rogersii*, although most specimens were not identified at the species level.

Genselia Knaus may have been a seed fern (Knaus, 1995; Taylor et al., 2009). This genus has bipinnate compound leaves, non-bifurcate rachises, and in some, bifurcations at the tip of stems, terminating in elongated sporangia (Knaus, 1994, 1995; Skog and Gensel, 1980; Taylor et al., 2009). It is found in the Lower Carboniferous Pocono and Price Formations of Pennsylvania, Maryland, West Virginia and Virginia, in the Appalachian Basin of North America. None of the *Genselia* specimens in our dataset was identified at the species level.

2.2. Paleoclimate and paleoenvironments

The locales from which fossil plants in this study were collected have a quasi-global distribution (Fig. 3A), although a large majority of the analyses ($n = 289$ of 309, or 93 % of the total dataset) are of specimens from the Appalachian Basin or other areas in eastern North America, at locales ranging from southwestern Virginia in the south to the Gaspé Peninsula of

Québec in the north (Fig. 3B). Laurentia (Paleozoic North America) drifted northward during the Devonian Period (416-359 Ma), causing the Appalachian Basin to shift from ~40–35 °S in the Early Devonian to ~30–25 °S in the Late Devonian (van der Voo, 1988, 1993). Thus, many of the present study locales moved from the humid temperate zone into the dry subtropical zone during this interval. However, this shift occurred at tectonic timescales (i.e., over a ~50-Myr interval) and was monotonic in direction (i.e., did not reverse), so any regional climate changes induced by plate motion during this interval are likely to have been slow and steady. More rapid climate changes (i.e., at timescales of a few million years or less) are likely to have had non-plate-tectonic causes. Most of the remaining fossil plant specimens ($n = 20$, 7 % of total) are from a few sites in northwestern China (Xu et al., 2015; Zheng et al., 2016) and Venezuela (Berry, 1994; Berry and Edwards, 1996).

The depositional environments in which the 12 fossil plant taxa of the present study grew ranged from coastal deltas to upper floodplains, with the former being the most common environment. Coastal deltas are represented, for example, by most of the fossil-plant-bearing strata of the Emsian Battery Point Formation in Gaspé Bay, Quebec, and the more western portions of the Emsian-lower Eifelian Campbellton Formation in New Brunswick (Griffing et al., 2000; Hotton et al., 2001; Kennedy et al., 2012, 2013). Especially at Gaspé Bay, paleocurrent data document mainly seaward transport of sediment by rivers, and multistory sandstone bodies are interpreted as deposits of high-sinuosity river channels (Griffing et al., 2000). Closer to the coastline, bidirectional cross-bedding records tidal influence (Bridge, 2000; Griffing et al., 2000) and some trace fossils (e.g., *Diplocraterion*) may represent brackish conditions within estuaries (Lawrence, 1986). Deltaic processes are evidenced by a complex stratigraphic

architecture of levees and crevasse splays, freshwater lakes and marshes, lacustrine deltas, brackish marshes and interdistributary bays, and sandy and muddy tidal flats (Griffing et al., 2000).

Coastal wetlands apparently developed frequently during the late Givetian to early Frasnian, especially at times of relatively high sea level, which resulted in marine incursions into generally terrestrial environments (Baird and Brett, 2008). Multiple wetland subenvironments are represented by stacked dark meter-thick sandstones at Riverside Quarry, Gilboa (Stein et al., 2012), dark siltstones and shales at Blenheim, Gilboa (Banks et al., 1972), and lighter gray to yellowish-brown sandstones at Manorkill Falls, Gilboa (Driese et al., 1997). Similar environments developed in the Late Devonian, as in the Famennian Hampshire Formation at Elkins, West Virginia (Scheckler, 1986a). In these settings, trees and understory plants appear to have had limited root penetration. During the Late Devonian, wetland floral diversity was relatively limited: *Archaeopteris* and *Rhacophyton* were endemic, although other taxa (especially arborescent lycophytes) occupied peri-swamp habitats and were washed in during floods and storms (Scheckler, 1986a). By the Tournaisian, coastal swamp habitats were increasingly dominated by lycophytes (Scheckler, 1986b; Rygel et al., 2006; Gensel and Pigg, 2010).

Upper floodplain settings are represented by fine-grained, multistory sandstone bodies and red mudstones in fining-upward successions lacking any evidence of marine influence. These deposits represent wide, meandering upland river channels and overbank (floodplain) deposits that were subject to frequent wet-dry cycles, e.g., through seasonal flooding (Cressler, 1999, 2006; Cressler et al., 2010). Examples of this type of environment are found in the

Famennian Duncannon Member of the Catskill Formation at Red Hill, Pennsylvania. Taphonomic observations indicate that both fossil plants and animals lived close to their sites of deposition. Fossil plant remains are often found in reduced lenses of greenish-gray mudstone, probably representing floodplain pond deposits related to major channel avulsion events (Cressler, 1999, 2006; Cressler et al., 2010).

3. Methods

The study specimens consist of well-preserved compression fossils of whole plants or parts of plants (i.e., stems, branches, and/or leaves) that are intact or nearly so (Fig. 4). All specimens were inspected using a binocular microscope to verify the state of fossil preservation, and a subset was examined using scanning electron microscopy. We initially bulk macerated each plant fossil in acid baths, yielding bulk organic material that was undifferentiated by plant anatomical parts. These specimens were digested successively in hydrochloric acid (HCl) and hydrofluoric acid (HF), with three rinses in distilled water between dissolution steps. We subsequently sampled the plant fossils by peeling pieces of organic material from the surface of each compression fossil, permitting us to analyze specific anatomical parts of each plant (e.g., stems, leaves, spines, and sporangia). These specimens were treated individually with HCl and HF, as needed, to remove rock matrix adhering to the organic material. Organic fragments were picked out of the residue, rinsed in distilled water, dried, and inspected under a binocular microscope to verify the removal of all impurities.

Carbon isotopic analyses were performed using EA-IRMS systems at the Stable Isotope Research Facility (SIRF) of the Department of Earth and Atmospheric Sciences of Indiana

University, Bloomington (IU) and the Department of Earth and Environmental Sciences of the University of Texas at Arlington (UTA). Analytical protocols were similar in the two laboratories: powdered samples were weighed into silver capsules that were acidified repeatedly with 6% sulfurous acid (H_2SO_3) in order to remove carbonate phases. Samples were analyzed using a Costech 4010 elemental analyzer interfaced with a Thermo Finnigan Conflo III device to a Thermo Finnigan Delta Plus XP isotope ratio mass spectrometer (IRMS). All isotopic results are reported in per mille (‰) variation relative to VPDB. At UTA, the average standard deviation was 0.04 ‰ for both the USGS-24 standard ($\delta^{13}\text{C} = -16.049$ ‰) and unknowns. At IU, the average standard deviation was 0.06 ‰ for Acetanilides #1, #2, and #3, and Corn starch #1 (Schimmelmann et al., 2016) and unknowns. Interlaboratory calibration was undertaken by analyzing a common set of 12 samples at both IU and UTA. The co-analyzed samples yielded a correlation (r^2) of 0.997 with an offset of 0.21 ‰ between the two labs, which was corrected for in the integrated C-isotope dataset by averaging the difference (i.e., by reducing UTA values by 0.10 ‰ and increasing IU values by 0.10 ‰).

4. Results

4.1. General results

In this study, we generated a total of 309 analyses of plant $\delta^{13}\text{C}$ from 190 unique plant specimens (Table S1). The number of analyses per taxon varied considerably: the four most analyzed taxa were *Archaeopteris* (80 analyses of 51 unique specimens), *Drepanophycus* (40/21), *Rhacophyton* (33/18), and *Tetraxylopteris* (31/21) (Table 1). About half of the specimens were analyzed twice, mostly to generate paired $\delta^{13}\text{C}$ values for different anatomical

parts (e.g., stems and leaves) but sometimes to check on the reproducibility of results for the same anatomical part of a single specimen. A handful of specimens were analyzed three or more times. The full dataset of 309 analyses yielded a mean $\delta^{13}\text{C}_p$ of -25.5‰ with a standard deviation of 1.8‰ and a range from -20.3‰ to -30.5‰ . Individual taxa yielded mean $\delta^{13}\text{C}_p$ ranging from $-22.3 \pm 1.3\text{‰}$ for *Haskinsia* to $-27.3 \pm 1.7\text{‰}$ for *Sawdonia* (Table 1).

The full $\delta^{13}\text{C}_p$ dataset exhibits a distinct secular pattern (as reflected in a LOWESS curve) that is dominated by a long-term shift toward higher $\delta^{13}\text{C}_p$ values through most of the Devonian and Early Carboniferous (note: this secular pattern is not shown and will not be considered here because it is the focus of a companion paper that is in preparation). To determine the non-secular component of total $\delta^{13}\text{C}_p$ variation, we calculated the difference between each $\delta^{13}\text{C}_p$ value and the age-equivalent mean for the full $\delta^{13}\text{C}_p$ dataset as given by the LOWESS curve, i.e., $\Delta^{13}\text{C}_p$ (Fig. 5). Thus, a fossil specimen with a $\Delta^{13}\text{C}_p$ of 0‰ has a carbon-isotopic composition identical to the LOWESS mean for the $\delta^{13}\text{C}_p$ dataset at a given time, and positive and negative $\Delta^{13}\text{C}_p$ values represent ^{13}C enrichment and depletion of the specimen, respectively, relative to the age-equivalent LOWESS mean value. In this manner, the overall tendency of each fossil plant taxon toward ^{13}C enrichment or depletion can be evaluated independently of long-term secular variation in the $\delta^{13}\text{C}_p$ dataset (Fig. 6). The non-secular variance in our $\delta^{13}\text{C}_p$ dataset has two components: (1) variance among different plant taxa ('intergeneric variation'), and (2) variance within individual taxa ('intrageneric variation'). The proportion of total variance in the $\Delta^{13}\text{C}_p$ dataset represented by intrageneric variance (i.e., $\sigma^2_{\text{intra}} / \sigma^2_{\text{total}}$) was calculated as:

$$\left(\sum_{i=1}^{12} [\Sigma(X-\mu_i)^2 / N_i] \right) / [\Sigma(X-\mu_t)^2 / N_t] \quad (1)$$

where X is the $\Delta^{13}\text{C}_p$ for a given fossil specimen, μ_i is the mean $\Delta^{13}\text{C}_p$ for a given plant genus i , and μ_t is the mean $\Delta^{13}\text{C}_p$ for all fossil specimens. The numerator of the equation sums the variances of the 12 fossil plant genera ($\sum_{i=1}^{12}$) calculated separately (i.e., $[\sum(X-\mu_i)^2 / N_i]$), and it is divided by the total variance for all fossil specimens (i.e., $[\sum(X-\mu_t)^2 / N_t]$) to yield the fraction of total variance attributable to intrageneric variance (70%); the remainder (30 %) represents intergeneric variance. Below, we analyze intrageneric and intergeneric patterns of variation as a function of possible taxonomic, anatomical, geographic, and environmental controls (Tables 2-3). Note that this analysis is conducted on the basis of $\Delta^{13}\text{C}_p$ values (Table S1), i.e., the deviations from the long-term secular $\delta^{13}\text{C}_p$ trend (Fig. 6), and is thus independent of the secular component of variance in the $\delta^{13}\text{C}_p$ dataset.

We considered the possibility of a geographic gradient in $\Delta^{13}\text{C}_p$, e.g., along a paleolatitudinal transect. Modern plants show a distinct latitudinal gradient in $\delta^{13}\text{C}_p$ that is a function of temperature and precipitation influences (Diefendorf et al., 2010) and/or light levels (Kohn, 2010). However, the high proportion (>90 %) of specimens from the Appalachian Basin in this study (Figs. 2-3) leads to a clustering of data within a relatively narrow range of paleolatitudes, resulting in our inability to identify any significant paleolatitudinal trends. A more robust test of geographic influences on Devonian fossil plant $\delta^{13}\text{C}$ will require a larger dataset covering a wider range of paleolatitudes.

4.2. Intrageneric variation in plant $\delta^{13}\text{C}$

For a subset of specimens ($n = 32$), we analyzed carbon from both the stem and the leaf or spine (not interpreted as a leaf homologue) of the fossil plant, allowing us to evaluate

variation in $\delta^{13}\text{C}_p$ as a function of plant anatomy. The assessment of anatomical variation was made for 6 plant genera (Table 3). Among these genera, four showed higher $\delta^{13}\text{C}_p$ for leaves/spines than for stems, although the difference was statistically significant (based on a Student's t-test) only for *Sawdonia*, which yielded the largest $\Delta^{13}\text{C}_{(\text{spine-stem})}$ value of any genus (1.6 ‰). The relatively small number of analyses per genus was a factor limiting the statistical significance of these results for individual plant taxa. All 6 genera together ($n = 32$) yielded a mean leaf/spine-vs-stem difference of 0.45 ‰, which is statistically significant ($p(\alpha) = 0.01$; Table 3). These results suggest that fossil plant leaves and spines are systematically slightly enriched in ^{13}C relative to stems of the same fossil plant specimen (as seen in 29 out of 32 of the paired analyses).

4.3. Intergeneric variation in plant $\delta^{13}\text{C}$

The 12 fossil plant genera of this study exhibit systematic differences in $\Delta^{13}\text{C}_p$ (Fig. 7). Three taxa yielded mean $\Delta^{13}\text{C}_p$ values that are significantly higher than the age-equivalent LOWESS mean: *Haskinsia* ($+2.4 \pm 1.5$ ‰; $n = 4$), *Leclercqia* ($+1.8 \pm 1.0$ ‰; $n = 24$), and *Psilophyton* ($+0.9 \pm 1.5$ ‰; $n = 25$), and one taxon yielded a significantly lower mean value: *Sawdonia* (-1.2 ± 1.8 ‰; $n = 28$) (note: for all, $p(\alpha) \leq 0.01$; Student's t-test; Table 2). The remaining eight taxa have mean $\Delta^{13}\text{C}_p$ values within ± 0.5 ‰ of the average for the dataset as a whole, among which only *Archaeopteris* yielded a significantly different mean (-0.4 ± 1.2 ‰; $p(\alpha) = 0.002$), mainly as a result of the exceptionally large number of analyses of this taxon ($n = 80$; Table 2). However, the result for *Archaeopteris* must be viewed cautiously because there are relatively few coeval Late Devonian taxa in our dataset from which to construct the LOWESS mean trend.

Thus, we consider only *Haskinsia*, *Leclercqia*, *Psilophyton*, and *Sawdonia* to deviate significantly in $\Delta^{13}\text{C}_p$ from age-equivalent LOWESS mean values.

The critical time interval of our dataset for evaluating interspecific $\Delta^{13}\text{C}_p$ variation is the late Early through late Middle Devonian (~405-385 Ma), during which 6 of the 12 plant taxa analyzed in this study have largely overlapping stratigraphic ranges. The overlap in stratigraphic ranges facilitates intergeneric comparisons, and, not coincidentally, all four of the taxa showing large deviations from mean LOWESS $\delta^{13}\text{C}_p$ values are present in this interval, with three taxa showing ^{13}C -enriched values (*Haskinsia*, *Leclercqia*, and *Psilophyton*) and one showing ^{13}C -depleted values (*Sawdonia*) (see above). Importantly, the offsets in $\delta^{13}\text{C}_p$ between these taxa are relatively constant despite an overall increase in mean LOWESS $\delta^{13}\text{C}_p$ of ~3 ‰ from the late Early Devonian to the late Middle Devonian (Algeo et al., in preparation). We attribute these taxon-specific differences in ^{13}C enrichment to local environmental controls (see Section 5.3). In contrast, the 6 plant genera analyzed from the early Late Devonian to Early Carboniferous (~385-345 Ma) have only partially overlapping stratigraphic ranges, with no more than 2 taxa present in any given time window. The more limited dataset for this time interval renders uncertain evaluation of deviations in $\Delta^{13}\text{C}_p$ from the mean LOWESS trend (thus our caution with regard to interpreting *Archaeopteris* $\Delta^{13}\text{C}_p$ values).

5. Discussion

5.1. Robustness of plant $\delta^{13}\text{C}$ proxy

The fossil plant specimens used in this project were preserved as compressions of stems, branches, leaves and spines (Fig. 4A-B). During the carbonization process, most water

416 and volatile organic compounds in the plant tissue were lost, and the remaining material was
417 preserved as a black organic film that consists largely of carbon (Guo et al., 2010). Even though
418 compressed as flat layers, many fossils show well-preserved cell-scale anatomical features such
419 as tracheids (Fig. 4C) and cuticle (Fig. 4D). These observations confirm that pre-burial bacterial
420 decay of these relatively resistant tissues was limited in the specimens that were selected for
421 carbon-isotopic analysis in this study.

422 Vascular land plants exhibit internal carbon isotopic variation, commonly with a >5 ‰
423 range in $\delta^{13}\text{C}$ among their constituent compounds (O'Leary, 1988; Pate and Arthur, 1998).
424 Although most types of compounds decay rapidly in the burial environment, the surviving
425 compounds generally retain their characteristic carbon isotopic compositions (Benner et al.,
426 1987). Cellulose and lignin are among the most resistant compound types and commonly
427 dominate the terrestrial organic fraction preserved in sedimentary rocks (Arens et al., 2000),
428 especially in compression fossils of the type analyzed in this study. Cellulose is up to ~3 ‰
429 heavier and lignin up to ~4 ‰ lighter than bulk plant $\delta^{13}\text{C}$ (Benner et al., 1987; O'Leary, 1988;
430 Marino and McElroy, 1991; Badeck et al., 2005). Biomarker analysis of plant compression fossils
431 typically yields a range of alkanes, alkenes, benzenes, pristenes, and polysaccharides derived in
432 large part from decay of these primary compounds (e.g., Gupta et al., 2007), so a part of the
433 variance in the $\Delta\delta^{13}\text{C}_p$ of an individual taxon in our dataset may reflect specimen-specific
434 variation in the proportions of cellulose versus lignin and their derivatives (Arens et al., 2000).

435 Burial diagenesis can potentially modify the $\delta^{13}\text{C}$ composition of organic material
436 through thermal cracking and release of methane at high temperatures (i.e., >160 °C, or beyond
437 the 'oil window'; Faure and Mensing, 2005). The released methane is depleted in ^{13}C by 10-

30 ‰, resulting in a small increase in the $\delta^{13}\text{C}$ of the remaining kerogen (generally <1 ‰). The thermal maturity of Devonian sediments in the central Appalachian Basin exhibits a strong lateral gradient: maturity is high on the eastern basin margin ($R_o > 2.0$; $T > 160^\circ\text{C}$) but low to moderate on the western basin margin ($R_o < 1.0$; $T < 100^\circ\text{C}$) (Obermajer et al., 1997; Milici and Swezey, 2006; Rowan, 2006; Repetski et al., 2008). In our study areas in eastern New York and central West Virginia, the maximum burial depths of Upper Devonian strata were from ~4 to 10 km, with peak burial temperatures estimated to have been between ~150 and 170 °C (Friedman and Sanders, 1982; Dorobek, 1989; Repetski et al., 2008). Thermal maturity is also variable in New Brunswick and the Gaspé Peninsula of Québec in eastern Canada. There, some Devonian strata have been buried just a few kilometers (Heroux et al., 1979), whereas other units have experienced maximum burial depths up to 12 km (Utting and Hamblin, 1991), yielding peak burial temperature estimates ranging from ~50 to 280 °C (Bertrand and Malo, 2001; Chi et al., 2001). We tested the possibility of burial thermal effects on our dataset by plotting $\delta^{13}\text{C}_p$ against inferred maximum burial depths, but this yielded almost no correlation ($r = +0.03$; $n = 309$; $p(\alpha) > 0.5$). We are cognizant that the carbon isotopic compositions of our plant fossils may have been influenced by bacterial and thermal decay. However, the $\delta^{13}\text{C}_p$ values of a single taxon at a specific outcrop tend to be relatively consistent in our dataset (mostly showing <1 ‰ variation), indicating a general lack of random diagenetic effects on the plant fossils during burial.

5.2. Sources of anatomical variation in plant $\delta^{13}\text{C}$

Among modern plants, different plant parts exhibit systematic differences in carbon isotopic compositions. Based on a compilation of >400 plant taxa, leaves are on average 0.96 ‰ and 1.91 ‰ more depleted in ^{13}C than roots and woody stems, respectively (Badeck et al., 2005). This pattern is strong but not universal as ~10 % of the taxa examined in that study yielded leaf $\delta^{13}\text{C}$ greater than stem $\delta^{13}\text{C}$. This pattern appears to be independent of climatic conditions, as plants from semi-arid habitats also exhibit leaf $\delta^{13}\text{C}$ lower than stem $\delta^{13}\text{C}$ (by an average of 1.51 ± 0.42 ‰) (Nilson and Sharifi, 1997). The origin of these differences is not known with certainty: two leading ideas relate it to carbon-isotopic fractionation in generating different types of organic compounds (see Section 5.1), or preferential release of ^{13}C -enriched respiratory CO_2 from leaves (Badeck et al., 2005).

In our dataset, leaves and spines are somewhat ^{13}C -enriched relative to stems (by 0.45 ‰ on average; Table 3; see Section 4.2). If the $\delta^{13}\text{C}$ values of leaves and spines were originally lower than that of stems by ~1-2 ‰ in Devonian plants, then burial decay must have caused a relative shift of +1.5 to +2.5 ‰ in leaf/spine $\delta^{13}\text{C}$ relative to stem $\delta^{13}\text{C}$. Such a shift might have occurred owing to a greater loss of lipids and/or increase in cellulose-derived compounds in the carbonized residue of leaves and spines relative to stems during the fossilization process. Further investigation will be needed to validate the existence and significance of internal $\delta^{13}\text{C}$ variation in Devonian and Carboniferous fossil plants.

5.3. Sources of intergeneric variance in plant $\delta^{13}\text{C}$

5.3.1. Water-use efficiency

One mechanism that can lead to systematic differences in $\delta^{13}\text{C}$ values between plant taxa is differences in water-use efficiency (WUE): plants that grow in wet habitats do not need to limit water loss and, hence, tend to open their stomata more widely in order to maximize CO_2 uptake at the cost of low water-use efficiency. This process leads to maximum internal-leaf $p\text{CO}_2$ and maximum carbon isotope discrimination (i.e., lower $\delta^{13}\text{C}_p$) (Farquhar et al., 1989). In contrast, plants that grow in drier habitats are forced to limit water loss through their stomata (i.e., higher water-use efficiency), leading to less CO_2 uptake, lower internal-leaf $p\text{CO}_2$, and reduced carbon-isotope discrimination (i.e., higher $\delta^{13}\text{C}_p$) as a result of a larger fraction of internal-leaf CO_2 being utilized in photosynthesis (Farquhar et al., 1989).

The carbon isotopic composition of modern C_3 land plants exhibits a strong negative relationship to mean annual precipitation (MAP) (Miller et al., 2001; Liu et al., 2005; Roden et al., 2005; Diefendorf et al., 2010; Kohn, 2010). An increase in MAP from 0 to 1000 mm yr^{-1} correlates with a ~4-5 ‰ depletion in $\delta^{13}\text{C}_p$, although this relationship is non-linear because $\delta^{13}\text{C}_p$ becomes less sensitive to changes in precipitation at high MAP (Diefendorf et al., 2010). This relationship exists because land plants are more commonly water-limited than carbon-limited and have a strong incentive to engage in greater water-use efficiency where water is scarce (Bacon, 2004; Macfarlane et al., 2004). Plants tend to reduce stomatal conductance in arid regions in order to conserve water, which causes a larger proportion of the CO_2 that diffuses into the leaf to be used in photosynthesis, resulting in reduced fractionation (relative to the atmospheric carbon source) and heavier $\delta^{13}\text{C}_p$ values (Farquhar et al., 1989; Brugnoli and Lauteri, 1991; Poss et al., 2000). The opposite pattern is found in humid regions, causing $\delta^{13}\text{C}_p$

values to become lighter. This relationship accounts for ~55 % of $\delta^{13}\text{C}$ variance among modern C_3 plants (Diefendorf et al., 2010).

A fundamental divide exists between species with high hydraulic conductivity and low embolism resistance and those with low hydraulic conductivity and high embolism resistance (Wilson, 2016). Mapping of hydraulic ecospace shows that the high-conductivity/low-safety-margin domain was occupied by many early vascular land plants, including trimerophytes, lycophytes, cladoxylopsids, and sphenopsids, all of which developed tracheids with scalariform pits that maximized hydraulic conductivity (Wilson and Knoll, 2010; Wilson and Fischer, 2011; Wilson, 2013, 2016). For some plants, this strategy was associated with perennially wet habitats in which the danger of desiccation was limited, e.g., as for arborescent lycophytes and other plants that inhabited tropical lowland swamps during the Carboniferous Period. For other plants, this hydraulic mechanism was associated with an ecological strategy based on rapid growth and generational overturn (known as ‘live fast, die young’), which possibly included all of the zosterophyllophytes and basal euphyllophytes of the present study. For example, *Psilophyton dawsonii* may have grown rapidly, as suggested by xylem with large pits similar to that of some modern small ferns whose aerial vegetative structures rarely persist for more than a year (Wilson, 2016). An additional factor in the high conductivity of early vascular plants was high atmospheric CO_2 levels, which permitted smaller and less numerous stomata (thus minimizing some types of cavitation hazards) and generally higher water-use efficiency (Sperry, 2003).

It has been suggested that early land plants were mostly generalists, occupying a wide range of habitats (Spicer, 1989; Meyer-Berthaud et al., 2003). However, the systematic

variation in $\Delta^{13}\text{C}_\text{p}$ among Devonian plant taxa observed in this study (Fig. 7), if due to environmental controls, implies that some early vascular land plants may have been more specific in their habitat preferences than previously thought. Wetter habitats may have been favored by most zosterophyllophytes, which typically had small, shallow root systems (Gensel et al., 2001; Xue, 2012). *Sawdonia* exhibits the relatively most ^{13}C -depleted compositions among the 12 study taxa (Fig. 7). It occupied a wide range of environments but was most common along the margins of interdistributary basins in lower delta plain settings (Griffing et al., 2000; Hotton et al., 2001; Kennedy et al., 2012). Sedimentologic investigations have not established for certain whether these basins were inundated mainly with brackish (Hotton et al., 2001) or fresh waters (Kennedy et al., 2012). The low $\Delta^{13}\text{C}_\text{p}$ compositions of *Sawdonia* documented in the present study are more consistent with fresh waters, because C_3 plants growing in coastal salt marshes or in areas of elevated groundwater salinity tend to become ^{13}C -enriched rather than ^{13}C -depleted (Brugnoli and Lauteri, 1991; Malamud-Roam and Ingram, 2001).

Drier habitats may have been occupied by some of the plant taxa examined in this study. In particular, *Leclercqia*, *Haskinsia*, and, to a lesser degree, *Psilophyton* exhibit ^{13}C -enriched compositions consistent with higher water-use efficiency (Fig. 7). This inference is supported by the anatomical features of these plants. The morphologies of *Leclercqia* and *Haskinsia* are similar to modern *Lycopodium* s.l., which often grows in seasonally dry habitats (Fernandez et al., 2008). In these taxa, densely crowded and overlapping leaves can create a boundary layer effect that is effective at minimizing water loss during gas exchange, and which is therefore favored in dry habitats. In addition, the leaves of *Leclercqia* and *Haskinsia* partially

enclosed the sporangia, protecting them from drying out (Meyer-Berthaud et al., 2003; Gensel and Kasper, 2005; Gensel and Albright, 2006). Certain features in *Psilophyton* may have adapted it to sunny, seasonally dry habitats, e.g., forked cylindrical lateral branchlets and a prominent outer cortex of axial fibers that provided mechanical support (possibly against wilting) and UV protection (via the high lignin content of fibrous cell walls) (Trant and Gensel, 1985; Gerrienne, 1995, 1997). These ^{13}C -enriched taxa may have grown where soils were better drained, e.g., on river levees or in upper floodplain settings that were further inland and somewhat drier than coastal environments, as proposed for *Psilophyton* (Hotton et al., 2001).

In areas of limited precipitation, increases in soil salinity can exert a strong influence on $\delta^{13}\text{C}_\text{p}$. Experimental and in-situ studies of modern C_3 plants have shown that $\delta^{13}\text{C}_\text{p}$ increases by 2-5 ‰ with rising soil salinity (Brugnoli and Lauteri, 1991; Poss et al., 2000; Winter and Holtum, 2005). A similar effect is observed in coastal salt marshes, where the $\delta^{13}\text{C}_\text{p}$ of C_3 marsh plants increases by several per mille with rising watermass salinity (Malamud-Roam and Ingram, 2001; Cloern et al., 2002). This effect is linked to the need to conserve water in more saline environments, leading to reduced stomatal conductance and photosynthetic fractionation and, thus, higher $\delta^{13}\text{C}_\text{p}$ values (Farquhar et al., 1989; Brugnoli and Lauteri, 1991; Poss et al., 2000). However, there is no evidence that any early vascular land plant grew in saline soils or brackish-water environments (Kennedy et al., 2012), and our results are consistent with this inference. The apparent water-use efficiencies of our fossil plant taxa suggested by their $\Delta^{13}\text{C}_\text{p}$ values (Fig. 7) correspond well with predictions of habitat preference based on paleoecological analyses (see Section 5.4). Further carbon isotopic study of early vascular land plants may prove invaluable in helping to define their habitat preferences.

5.3.2. Forest understory $\delta^{13}\text{C}$ gradient

A second environmental factor that can lead to systematic differences in $\delta^{13}\text{C}$ values between plant taxa is vertical $\delta^{13}\text{C}_{\text{CO}_2}$ variation in the forest understory atmosphere. Forests are characterized by a higher rate of soil respiration than most other vegetation biomes (Raich and Tufekciogul, 2000), as well as by more limited air mass exchange than in open landscapes owing to the sheltering effects of a closed canopy (Feigenwinter et al., 2004). This combination of factors results in elevated $p\text{CO}_2$ in the forest understory atmosphere along with a significant vertical $\delta^{13}\text{C}_{\text{CO}_2}$ gradient (Jackson et al., 1993). At 0.5 m above the forest floor, up to ~20 % of CO_2 is soil-derived (Da Silveira et al., 1989), exposing low-growing plants to strongly ^{13}C -depleted (–28 to –25 ‰) soil-respired CO_2 (Farquhar et al., 1989; Jackson et al., 1993). As a consequence, CO_2 in the forest understory atmosphere may be up to ~3-5 ‰ depleted relative to CO_2 in the open atmosphere, and this isotopic signature can be transferred to understory plants (Schleser and Jayasekera, 1985; Flanagan et al., 1996). In modern forests, >50 % of $\delta^{13}\text{C}$ variance in understory plants is linked to local variations in $\delta^{13}\text{C}_{\text{CO}_2}$ (Da Silveira et al., 1989).

‘Canopy effects’ are unlikely to have played any role in the intergeneric differences in $\Delta^{13}\text{C}_\text{p}$ of the present study. The largest differences in $\Delta^{13}\text{C}_\text{p}$ values are seen among Early and early Middle Devonian plants that existed prior to the first forests (Fig. 6). These early plants grew in patchy floral communities consisting largely of clonal vegetation whose shoots would have lacked any capacity to limit air mass exchange (Gensel and Edwards, 2001; Edwards and Richardson, 2004), and whose limited productivity did not generate high excess soil $p\text{CO}_2$ (Mora et al., 1996; Elick et al., 1998). The appearance of the first forests during the late Givetian to

early Frasnian (Decombeix et al., 2011; Stein et al., 2012; Berry and Marshall, 2015), an event known as 'afforestation' (Scheckler, 2001), resulted in denser floral communities with closed canopies formed by large leafy trees that, for the first time, would have generated understory niches with varying environmental characteristics. In our dataset, the only co-existing Late Devonian taxa that have the potential to exhibit $\Delta^{13}\text{C}_\text{p}$ differences due to a 'canopy effect' are *Archaeopteris*, a canopy tree, and *Rhacophyton*, a much smaller woody bush (Fig. 1). However, *Rhacophyton* yields $\Delta^{13}\text{C}_\text{p}$ values that are on average $\sim 1.5\text{‰}$ heavier than those of *Archaeopteris* (Fig. 6), which is the opposite of the pattern expected for canopy-versus-understory plants. We infer that the higher $\Delta^{13}\text{C}_\text{p}$ values of *Rhacophyton* are more likely due to its preference for dry habitats, leading to enhanced water-use efficiency (see Section 5.3.1).

5.4. Relationship of $\delta^{13}\text{C}$ to habitat preferences of early vascular plants

Although it has been proposed that early land plants were mostly generalists (Spicer, 1989; Meyer-Berthaud et al., 2003), the carbon-isotopic evidence of the present study suggests that Devonian and Early Carboniferous plant taxa varied considerably in their habitat preferences. Systematic interspecific differences in $\Delta^{13}\text{C}_\text{p}$ among the 12 taxa examined (Fig. 7) were probably controlled mainly by water-use efficiency, related to humidity levels at their sites of growth (see Section 5.3.1). To illustrate inferred differences in habitat, we have generated reconstructions of Early, Middle, and Late Devonian terrestrial ecosystems (Fig. 8A-C). These reconstructions focus on the distributions of the 12 plant genera of the present study, with the spatial distribution of each taxon shown to conform to its relative water-use efficiency as inferred from taxon-specific differences in $\Delta^{13}\text{C}_\text{p}$.

Our reconstructions of Devonian land-plant habitats were informed by the findings of earlier studies of terrestrial ecosystems. One consideration in habitat reconstruction is the in-situ versus transported character of land plant fossils. Most plant fossils have been transported prior to final deposition and burial (e.g., [Pratt and van Heerde, 2017](#)), although fossils that consist of intact stems and leaves and are well-preserved suggest limited transport distances ([Hotton et al., 2001](#); [Allen and Gastaldo, 2006](#); [Kennedy et al., 2012](#); [Gastaldo, 2016](#)). Nearly all of the fossil plant specimens of the present study are well-preserved and relatively intact, often consisting of complete stems, twigs or leaves, and more rarely of dense intertangled mats of vegetation. Thus, it is unlikely that they were transported long distances prior to burial (cf. [Berry and Edwards, 1996](#); [Gastaldo, 2016](#); [Pratt and van Heerde, 2017](#)). Such fossils are considered to be 'parautochthonous', i.e., transported to only a limited degree and thus useful for environmental interpretations of their growth habitat ([Hotton et al., 2001](#); [Allen and Gastaldo, 2006](#); [Kennedy et al., 2012](#)).

During the Early to Middle Devonian, land plants were concentrated in coastal delta and lower floodplain habitats ([Hotton et al., 2001](#)). From the Middle Devonian, land plants began to penetrate more deeply into continental interiors and increasingly occupied upper floodplain settings ([Algeo et al., 1995](#); [Berry et al., 2000](#); [Berry and Fairon-Demaret, 2001](#); [Cressler et al., 2010](#); [Retallack and Huang, 2011](#)). However, the upland regions of continental interiors remained mostly devoid of vegetation until the advent of seed plants in the latest Devonian to Early Carboniferous ([Algeo and Scheckler, 1998](#); [Decombeix et al., 2011](#)). Early to Middle Devonian plants appear to exhibit a wider range of $\Delta\delta^{13}\text{C}_\text{p}$ variation than Late Devonian to Early Carboniferous plants. This pattern may reflect greater environmental variation in Early-Middle

Devonian terrestrial habitats. During this early stage of landscape colonization, higher land plants had probably densely colonized some areas (e.g., delta plains) and sparsely colonized others (e.g., upland floodplains), while leaving broad continental interiors largely unvegetated. As a consequence of this mosaic floral distribution, land areas may have exhibited substantial spatial variation in environmental characteristics (e.g., humidity, soil development, albedo, etc.) (cf. [Edwards and Richardson, 2004](#)).

Early Devonian terrestrial ecosystems ([Fig. 8A](#)) have been analyzed in studies of the Emsian-age Battery Point Formation on the Gaspé Peninsula of Québec and the Campbellton Formation in New Brunswick. Facies A of the Cap-aux-Os Member of the Battery Point Formation contains unimodal paleocurrent indicators in coastal delta-plain facies that have been interpreted as crevasse splay or storm washover deposits ([Hotton et al., 2001](#)). The most common taxon in this setting is *Sawdonia*, which is inferred to have lived close to the shoreline on the margins of interdistributary basins that were subject to frequent flooding. [Hotton et al. \(2001\)](#) inferred that flooding events introduced brackish or marine waters, but [Kennedy et al. \(2012\)](#) inferred that similar occurrences of *Sawdonia* in the Campbellton Formation existed in fully freshwater habitats. Facies B of the Cap-aux-Os Member represents prograding fluvial channel deposits, hence mainly freshwater conditions, although some of the sandstones contain asymmetric bimodal cross-bedding indicative of tidal influence ([Hotton et al., 2001](#)). This facies contains abundant *Pertica* and *Drepanophycus* (as well as other taxa not analyzed in this study) that are thought to have grown along lower floodplain channel margins, whereas *Psilophyton* was inferred to have occupied an equivalent habitat somewhat further upstream. Similar assemblages containing *Pertica*, *Drepanophycus*, *Psilophyton*, and *Leclercqia* have been

reported from coastal delta plain and freshwater fluvial deposits of the Campbellton Formation in New Brunswick (Kennedy et al., 2012) and the Trout Valley Formation of Maine (Allen and Gastaldo, 2006).

Early Devonian vascular plants were dominantly ground-hugging rhizomatous lycophytes (*Sawdonia*, *Drepanophycus*, *Leclercqia*, *Haskinsia*) and shrubby trimerophytes (*Psilophyton*, *Pertica*), mostly with heights of <1 m (Fig. 1B; Algeo and Scheckler, 1998). Many Early Devonian plants grew in dense monospecific stands, facilitated by predominantly rhizomatous growth strategies, allowing local resource domination (Fig. 8A; DiMichele and Hook, 1992; Xue, 2012). However, mixtures of plant fossil debris in many formations imply that such patchiness was relatively local, and that there was considerable taxonomic heterogeneity in Early Devonian landscapes at a slightly larger spatial scale (Allen and Gastaldo, 2006; Cressler et al., 2010; Gastaldo, 2016). This pattern may reflect habitat specialization ('floral partitioning') on the basis of environmental variability in soil moisture, nutrient levels, or environmental characteristics such as frequency of flooding or other disturbances (Allen and Gastaldo, 2006; Greb et al., 2006).

By the late Middle Devonian (Givetian; Fig. 8B), the development of secondary supporting tissues resulted in an increase in average plant heights and the development of the earliest forests in coastal wetland regions (DiMichele and Hook, 1992; Greb et al., 2006; Mintz et al., 2010; Stein et al., 2012; Berry and Marshall, 2015). Shrubby stands of aneurophytalean progymnosperms, lycopsids, cladoxylopsids, and zosterophyllophytes were widespread in lowland areas (Fig. 8B; Berry and Fairon-Demaret, 2001; Xue et al., 2018). Early forests consisted of the cladoxylopid tree *Eospermatopteris* with an understory flora that included

lycopsids and aneurophytalean progymnosperm shrubs such as *Tetraxylopteris* (Driese et al., 1997; Stein et al., 2007, 2012; Mintz et al., 2010; Xu et al., 2017). The rapid global spread of forests may have been linked to archaeopteridalean progymnosperm trees with laminate leaves and deep root systems. The appearance of early members of this clade in the early Givetian is signaled by the microspore *Geminospora lemurata* and the megaspore *Contagisporites optivus* (Marshall, 1996; Turnau, 2014), and their widespread presence in mid to late Givetian landscapes is attested by specimens of *Svalbardia* or proto-*Archaeopteris* (Berry, 2008).

Late Devonian landscapes were dominated by dense monospecific stands of *Archaeopteris*, with *Rhacophyton* and other shrubby plants either occupying open terrain or present as understory elements in forests (Fig. 8C). Archaeopterid progymnosperms formed extensive forests in both lower and upper floodplain environments (Scheckler, 1986a; Beck and Wight, 1988; Meyer-Berthaud et al., 1999), and *Archaeopteris* is thought to have generally favored dry riparian habitats (Cressler, 1999, 2006; Retallack and Huang, 2011). *Rhacophyton* grew as a bushy understory plant in forests subject to frequent ground fires (Cressler, 2001, 2006), although it also flourished in peat wetlands (Greb et al., 2006). The earliest known seed plants became established in the mid-Famennian as small bushy colonizers in disturbed habitats, but they did not become important members of terrestrial ecosystems until the Carboniferous (Scheckler, 1986b; Streel et al., 2000; Decombeix et al., 2011).

5.5. Middle/Late Devonian floral turnover

Floral turnovers were a regular feature of early Devonian plant communities owing to the rapid pace of evolutionary innovation at that time (Cascales-Miñana et al., 2010). The rhyniophytes, which, as the earliest tracheophyte clade, were morphologically simple, peaked in dominance during the Late Silurian and earliest Devonian (Lochkovian). By the Early Devonian, the zosterophyllophytes had evolved lateral sporangia, pseudomonopodial branching, and a rhizomatous habit, establishing them as the most morphologically complex, diverse, and abundant clade (Gensel, 1992; Edwards and Richardson, 2004; Cascales-Miñana and Meyer-Berthaud, 2015). They peaked during the mid to late Early Devonian (Pragian-Emsian) but gradually declined as lycophytes and basal euphyllophytes gained in importance during the early Middle Devonian (Eifelian) (Hao and Xue, 2013b; Cascales-Miñana and Meyer-Berthaud, 2015). Among other innovations, basal euphyllophytes developed planate or laminate lateral branched systems resembling proto-leaves (Gerrienne et al., 2014). The turnovers among these clades appear to have been competitive replacements that were not clearly linked to any specific climatic or tectonic event (Cascales-Miñana and Meyer-Berthaud, 2015).

The Givetian/Frasnian (Middle/Late Devonian) transition appears to mark another major floral turnover (Raymond and Metz, 1995; Silvestro et al., 2015). During this event, primitive spore-bearing plants such as zosterophyllophytes experienced high extinction rates (Cascales-Miñana and Meyer-Berthaud, 2014) and euphyllophytes, including both lignophytes and non-lignophytes, underwent a major diversification (Xue et al., 2018). Among the taxa of the present study, the euphyllophytes *Psilophyton* and *Pertica* declined in the Emsian-Eifelian, and the zosterophyllophytes and lycopsids *Drepanophycus*, *Sawdonia*, *Leclercqia*, and *Haskinsia*

declined in the Givetian (Fig. 2). Although some of these taxa (e.g., *Drepanophycus*) persisted through the Famennian, their relative importance in the Late Devonian was greatly reduced. This floral turnover is even more apparent in palynomorph assemblages, which exhibit a sharp decline in diversity and rapid turnover at the Middle/Late Givetian boundary across Russia and eastern Europe and, to a lesser degree, in France, although it has not reported to date from North America (Obukhovskaya, 2000; Turnau, 2014).

One important aspect of the Givetian-Frasnian floral turnover was the appearance and diversification of arborescent (tree-sized) lycophytes and lignophytes (Wang et al., 2005; Galtier and Meyer-Berthaud, 2006; Meyer-Berthaud et al., 2010; Decombeix et al., 2011). In this study, the Givetian/Frasnian transition is marked by short-lived arborescent taxa such as *Wattieza* and *Tetraxylopteris*, after which longer-ranging arborescent taxa such as *Archaeopteris* became dominant in the Late Devonian (Fig. 2). The evolutionary development of trees led directly to the first forests, which were of Givetian (late Middle Devonian) to earliest Frasnian (early Late Devonian) age. They were dominated by tree-sized pseudosporochnalean cladoxylopsids such as *Wattieza/Eospermatopteris* with an understory that included aneurophytaleans such as *Tetraxylopteris*, as well as possibly herbaceous and arborescent lycopsids (Driese et al., 1997; Stein et al., 2007, 2012; Mintz et al., 2010). The known distribution of pseudosporochnalean forests is mostly limited to the Euramerican continental block, but they were succeeded by archaeopterid-dominated forests on all continents during the later Frasnian to Famennian (Scheckler, 1986a, 2001). The advent of forests probably generated selective pressures on terrestrial plants either to evolve larger size or to adapt to a shaded understory. The near-simultaneous appearance of woody supporting tissues and tree-sized stature among

arborescent lycophytes, sphenopsids, and lignophytes may imply the first strategy (Mosbrugger, 1990; Meyer-Berthaud et al., 2010). Less certain, due to difficulties in sampling, is what might be inferred about plant evolution in the understory. Nevertheless, the spread of forests likely had multifaceted, profound and irreversible consequences for both terrestrial floral communities and global climate conditions.

6. Conclusions

Carbon-isotopic variation among 12 genera of Devonian-Early Carboniferous vascular land plants provides insights into their habitat preferences and water-use efficiencies. A total of 309 analyses of 190 unique specimens yielded $\delta^{13}\text{C}_\text{p}$ values ranging from -20.3‰ to -30.5‰ with a mean of -25.5‰ , similar to the range and mean of $\delta^{13}\text{C}$ values for modern C_3 land plants. *Sawdonia* yielded the most ^{13}C -depleted values (mean $-27.1 \pm 1.7\text{‰}$; $n = 28$), reflecting lower water-use efficiency that was probably related to growth in wetter habitats such as lower delta plains. Because salt-marsh vegetation is typically ^{13}C -enriched, the strongly ^{13}C -depleted composition of *Sawdonia* is suggestive of growth in freshwater habitats. In contrast, *Leclercqia*, *Haskinsia*, and *Psilophyton* yielded relatively ^{13}C -enriched values (means $-23.0 \pm 1.6\text{‰}$, $-22.3 \pm 1.3\text{‰}$, and $-24.8 \pm 1.6\text{‰}$, respectively), reflecting higher water-use efficiency related to growth in drier habitats such as upper floodplains. This inference is supported by anatomical adaptations to drier conditions exhibited by these taxa, e.g., related to leaf arrangement and structure. A large majority of our specimens ($>90\%$) are from the Appalachian Basin of eastern North America, precluding effective assessment of large-scale geographic patterns of carbon-isotopic variation in these fossil plant taxa. This study

demonstrates that investigations of the carbon-isotopic composition of well-preserved plant fossils have the potential to yield insights regarding habitat preferences and ecosystem structure in paleofloral communities.

Acknowledgments

Thanks to the Peabody Museum of Yale University, the State Museum of New York, and the Smithsonian Institution for access to their paleobotanical collections. ZZW thanks Xinda Hu, Teng Teng, and N. Doug Rowe for field assistance. This study was supported by graduate research grants to ZZW from the American Association of Petroleum Geologists, the Geological Society of America, the Paleontological Society, the Society for Sedimentary Geology, and the University of Cincinnati Research Council.

References

- Algeo, T.J., Scheckler, S.E., 1998. Terrestrial-marine teleconnections in the Devonian: links between the evolution of land plants, weathering processes, and marine anoxic events. *Phil. Trans. Royal Society (B)* 353, 113-130.
- Algeo, T.J., Berner, R.A., Maynard, J.B., Scheckler, S.E., 1995. Late Devonian oceanic anoxic events and biotic crises: "rooted" in the evolution of vascular land plants? *GSA Today* 5(3), 45 & 64-66.
- Algeo, T.J., Scheckler, S.E., Maynard, J.B., 2001. Effects of early vascular land plants on weathering processes and global chemical fluxes during the Middle and Late Devonian. In:

786 Gensel, P., Edwards, D. (Eds.), *Plants Invade the Land: Evolutionary and Environmental*
787 *Perspectives*. Columbia Univ. Press, New York, pp. 213-236.

788 Allen, J.P., Gastaldo, R.A., 2006. Sedimentology and taphonomy of the Early to Middle Devonian
789 plant-bearing beds of the Trout Valley Formation, Maine. In: Greb, S.F., DiMichele, M.A.
790 (Eds.), *Wetlands Through Time*. Geol. Soc. Am. Spec. Publ. 399, pp. 57-78.

791 Andrews, H.N., Phillips, T.L., 1968. *Rhacophyton* from the Upper Devonian of West Virginia. Bot.
792 J. Linn. Soc. 61, 37-64.

793 Andrews, H.N., Kasper, A., Mencher, E., 1968. *Psilophyton forbesii*, a new Devonian plant from
794 northern Maine. Bulletin of the Torrey Botanical Club 95(1), 1-11.

795 Arens, N.C., Jahren, A.H., Amundson, R., 2000. Can C3 plants faithfully record the carbon
796 isotopic composition of atmospheric carbon dioxide? *Paleobiology* 26(1), 137-164.

797 Bacon, M.A., 2004. Water-use efficiency in plant biology. In: Bacon, M.A. (Ed), *Water-Use*
798 *Efficiency in Plant Biology*. Blackwell, Oxford, pp. 1-26.

799 Badeck, F.W., Tcherkez, G., Nogues, S., Piel, C., Ghashghaie, J., 2005. Post-photosynthetic
800 fractionation of stable carbon isotopes between plant organs—a widespread
801 phenomenon. *Rapid Comm. Mass Spectr.* 19(11), 1381-1391.

802 Baird, G.C., Brett, C.E., 2008. Late Givetian Taghanic bioevents in New York State: New
803 discoveries and questions. *Bulletin of Geosciences (Czech Geological Survey)* 83(4), 357-
804 370.

805 Banks, H.P., Bonamo, P.M., Grierson, J.D., 1972. *Leclercqia complexa* gen. et sp. nov., a new
806 lycopod from the Late Middle Devonian of eastern New York. *Rev. Palaeobot. Palynol.*
807 14(1-2), 19-40.

808 Banks, H.P., Leclercq, S.M, Hueber, F.M., 1975. Anatomy and morphology of *Psilophyton*
809 *dawsoni*, sp. n. from the Late Lower Devonian of Quebec (Gaspé) and Ontario, Canada.
810 *Palaeontogr. Amer.* 8(48), 77-127.

811 Beck, C.B., 1957. *Tetraxylopteris schmidtii* gen. et sp. nov., a probable pteridosperm precursor
812 from the Devonian of New York. *Am. J. Bot.* 42, 743-749.

813 Beck, C.B., 1962. Reconstruction of *Archaeopteris*, and further consideration of its phylogenetic
814 position. *Am. J. Bot.* 49(4), 373-382.

815 Beck, C.B., Wight, D.C., 1988. Progymnosperms. In: Beck, C.B. (Ed.), *Origin and Evolution of*
816 *Gymnosperms*. Columbia Univ. Press, New York, pp. 1-84.

817 Beerling, D.J., Lake, J.A., Berner, R.A., Hickey, L.J., Taylor, D.W., Royer, D.L., 2002. Carbon
818 isotope evidence implying high O₂/CO₂ ratios in the Permo-Carboniferous atmosphere.
819 *Geochim. Cosmochim. Acta* 66, 3757-3767.

820 Benca, J.P., Carlisle, M.H., Bergen, S., Strömberg, C.A., 2014. Applying morphometrics to early
821 land plant systematics: A new *Leclercqia* (Lycopsida) species from Washington State,
822 USA. *Am. J. Bot.* 101(3), 510-520.

823 Benner, R., Fogel, M.L., Sprague, E.K., Hodson, R.E., 1987. Depletion of ¹³C in lignin and its
824 implications for stable carbon isotope studies. *Nature* 329, 708-710.

825 Berner, R.A., 1997. The rise of plants and their effect on weathering and atmospheric CO₂.
826 *Science* 276(5312), 544-546.

827 Berry, C.M., 1994. First record of the Devonian lycophyte *Leclercqia* from South America.
828 *Geological Magazine* 131(2), 269-272.

829 Berry, C. M., 2008. The Middle Devonian plant collections of François Stockmans reconsidered.
830 Geologica Belgica 12, 25-30.

831 Berry, C.M., Edwards, D., 1996. The herbaceous lycophyte *Haskinsia* Grierson and Banks from
832 the Devonian of western Venezuela, with observations on the leaf morphology and fertile
833 specimens. Bot. J. Linn. Soc. 122, 103-122.

834 Berry, C.M., Fairon-Demaret, M., 2001. The Middle Devonian flora revisited. In: Gensel, P.,
835 Edwards, D. (Eds.), Plants Invade the Land: Evolutionary and Environmental Perspectives.
836 Columbia Univ. Press, New York, pp. 120-139.

837 Berry, C.M., Fairon-Demaret, M., 2002. The architecture of *Pseudosporochnus nodosus* Leclercq
838 et Banks: A Middle Devonian cladoxylopid from Belgium. Int. J. Plant Sci. 163, 699-713.

839 Berry, C.M., Marshall, J.E.A., 2015. Lycopod forests in the early Late Devonian paleoequatorial
840 zone of Svalbard. Geology 43, 1043-1046.

841 Berry, C.M., Morel, E., Mojica, J., Villarroel, C., 2000. Devonian plants from Colombia, with
842 discussion of their geological and palaeogeographical context. Geol. Mag. 137, 257-268.

843 Bertrand, R., Malo, M., 2001. Source rock analysis, thermal maturation, and hydrocarbon
844 generation in the Siluro-Devonian rocks of the Gaspé Belt basin, Canada. Bull. Can. Petr.
845 Geol. 49, 238-261.

846 Bonamo, P.M., Banks, H.P., 1967. *Tetraxylopteris schmidtii*: its fertile parts and its relationships
847 within the Aneurophytales. Am. J. Bot. 54(6), 755-768.

848 Bonamo, P.M., Banks, H.P., Grierson, J.D., 1988. *Leclercqia*, *Haskinsia*, and the role of leaves in
849 delineation of Devonian lycopod genera. Bot. Gaz. 149(2), 222-239.

850 Boyce, C.K., Hotton, C.L., Fogel, M.L., Cody, G.D., Hazen, R.M., Knoll, A.H., Heuber, F.M., 2007.
851 Devonian landscape heterogeneity recorded by a giant fungus. *Geology* 35, 399-402.

852 Bridge, J.S., 2000. The geometry, flow patterns and sedimentary processes of Devonian rivers
853 and coasts, New York and Pennsylvania, USA. In: Friend, P.F., Williams, B.P.J. (Eds.), *New*
854 *Perspectives on the Old Red Sandstone*. Geol. Soc. Lond. Spec. Publ. 180, pp. 85-108.

855 Brugnoli, E., Lauteri, M., 1991. Effects of salinity on stomatal conductance, photosynthetic
856 capacity, and carbon isotope discrimination of salt-tolerant (*Gossypium hirsutum* L.) and
857 salt-sensitive (*Phaseolus vulgaris* L.) C3 non-halophytes. *Plant Physiol.* 95, 628-635.

858 Cascales-Miñana, B., Meyer-Berthaud, B., 2014. Diversity dynamics of Zosterophyllopsida.
859 *Lethaia* 47(2), 205-215.

860 Cascales-Miñana, B., Meyer-Berthaud, B., 2015. Diversity patterns of the vascular plant group
861 Zosterophyllopsida in relation to Devonian paleogeography. *Palaeogeogr. Palaeoclimatol.*
862 *Palaeoecol.* 423, 53-61.

863 Cascales-Miñana, B., Muñoz-Bertomeu, J., Ros, R., Segura, J., 2010. Trends and patterns in the
864 evolution of vascular plants: macroevolutionary implications of a multilevel taxonomic
865 analyses. *Lethaia* 43, 545-557.

866 Chi, G., Lavoie, D., Bertrand, R., 2001. Hydrocarbon inclusions entrapped in diagenetic minerals
867 in Paleozoic basins in eastern Canada—Implications for hydrocarbon exploration. *Geol.*
868 *Surv. Canada Contr.* No. 2000252.

869 Cloern, J.E., Canuel, E.A., Harris, D., 2002. Stable carbon and nitrogen isotope composition of
870 aquatic and terrestrial plants of the San Francisco Bay estuarine system. *Limnol. Oceanogr.*
871 47(3), 713-729.

872 Cornet, B., Phillips, T.L., Andrews, H.N., 1977. The morphology and variation in *Rhacophyton*
873 *ceratangium* from the Upper Devonian and its bearing on frond evolution.
874 *Palaeontographica* (B) 158, 105-129.

875 Cressler, W.L., III, 1999. Site-analysis and floristics of the Late Devonian Red Hill locality,
876 Pennsylvania, an *Archaeopteris*-dominated plant community and early tetrapod site.
877 Unpubl. dissertation, U. of Pennsylvania, Philadelphia, Pennsylvania, 156 pp.

878 Cressler, W.L., III, 2001. Evidence of earliest known wildfires. *Palaos* 16, 171-174.

879 Cressler, W.L., III, 2006. Plant paleoecology of the Late Devonian Red Hill locality, north-central
880 Pennsylvania, an *Archaeopteris*-dominated wetland plant community and early tetrapod
881 site. In: Greb, S.F., DiMichele, M.A. (Eds.), *Wetlands Through Time*. Geol. Soc. Am. Spec.
882 Publ. 399, pp. 79-102.

883 Cressler, W.L., III, Daeschler, E.B., Slingerland, R., Peterson, D.A., 2010. Terrestrialization in the
884 Late Devonian: a palaeoecological overview of the Red Hill site, Pennsylvania, USA. In:
885 Vecoli, M., Clément, G., Meyer-Berthaud, B. (eds.), *The Terrestrialization Process:*
886 *Modelling Complex Interactions at the Biosphere-Geosphere Interface*, Geological Society
887 of London, Spec. Publ. 339, pp. 111-128.

888 Da Silveira Lobo Sternberg, L., Mulkey, S.S., Wright, S.J., 1989. Ecological interpretation of leaf
889 carbon isotope ratios: influence of respired carbon dioxide. *Ecology* 70(5), 1317-1324.

890 Decombeix, A.-L., Meyer-Berthaud, B., Galtier, J., 2011. Transitional changes in arborescent
891 lignophytes at the Devonian-Carboniferous boundary. *J. Geol. Soc. London* 168, 547-557.

892 Diefendorf, A.F., Mueller, K.E., Wing, S.L., Freeman, K.H., 2010. Global patterns in leaf ¹³C
893 discrimination and implications for studies of past and future climate. Proc. Nat. Acad. Sci.
894 USA 107, 5738-5743.

895 DiMichele, W.A., Hook, R.W., 1992. Paleozoic terrestrial ecosystems. In: Behrensmeyer, A.K.,
896 Damuth, J.D., DiMichele, W.A., Potts, R., Sues, H.-D., Wing, S.L. (Eds.), Terrestrial
897 Ecosystems Through Time. Univ. Chicago Press, Chicago, pp. 205-325.

898 Dittrich, H.S., Matten, L.C., Phillips, T.L., 1983. Anatomy of *Rhacophyton ceratangium* from the
899 Upper Devonian (Famennian) of West Virginia. Rev. Palaeobot. Palynol. 40, 127-47.

900 Doran, J.B. 1980. A new species of *Psilophyton* from the Lower Devonian of northern New
901 Brunswick, Canada. Can. J. Bot. 58, 2241-2262.

902 Doran, J.B., Gensel, P.G., Andrews, H.N., 1978. New occurrences of trimerophytes from the
903 Devonian of eastern Canada. Can. J. Bot. 56, 3052-3068.

904 Dorobek, S., 1989. Migration of orogenic fluids through the Siluro-Devonian Helderberg Group
905 during late Paleozoic deformation: constraints on fluid sources and implications for thermal
906 histories of sedimentary basins. Tectonophysics 159, 25-45.

907 Driese, S.G., Mora, C.I., Elick, J.M., 1997. Morphology and taphonomy of root and stump casts
908 of the earliest trees (Middle to Late Devonian), Pennsylvania and New York, USA. Palaios
909 12, 524-537.

910 Edwards, D., 2003. Xylem in early tracheophytes. Plant Cell and Environment 26, 57-72.

911 Edwards, D., Richardson, J.B., 2004. Silurian and Lower Devonian plant assemblages from the
912 Anglo-Welsh Basin: A palaeobotanical and palynological synthesis. Geol. J. 39(3-4), 375-
913 402.

914 Elick, J.M., Driese, S.G., Mora, C.I., 1998. Very large plant and root traces from the Early to
 915 Middle Devonian: Implications for early terrestrial ecosystems and atmospheric $p\text{CO}_2$.
 916 *Geology* 26(2), 143-146.

917 Farquhar, G.D., O'Leary, M.H., Berry, J.A., 1982. On the relationship between carbon isotope
 918 discrimination and the intercellular carbon dioxide concentration in leaves. *Austral. J. Plant*
 919 *Phys.* 9, 121-137.

920 Farquhar, G.D., Ehleringer, J.R., Hubick, K.T., 1989. Carbon isotope discrimination and
 921 photosynthesis. *Ann. Rev. Plant Phys. Plant Mol. Biol.* 40, 503-537.

922 Faure, G., Mensing, T.M., 2005. *Isotopes: Principles and Applications*, 3rd ed. John Wiley & Sons,
 923 Hoboken, New Jersey, 897 pp.

924 Feigenwinter, C., Bernhofer, C., Vogt, R., 2004. The influence of advection on the short term
 925 CO_2 -budget in and above a forest canopy. *Boundary-Layer Meteorology* 113(2), 201-224.

926 Feng, J., Xu, H., Jiang, Q., 2014. Study on the stable carbon isotope of the Mid-Late Devonian
 927 plants from north Xinjiang. *Bot. Res.* 3, 227-237 (in Chinese with English abstract).

928 Fernandez, N., Messuti, M.I., Fontenla, S., 2008. Arbuscular mycorrhizas and dark septate fungi
 929 in *Lycopodium paniculatum* (Lycopodiaceae) and *Equisetum bogotense* (Equisetaceae) in a
 930 valdivian temperate forest of Patagonia, Argentina. *Am. Fern J.* 98(3), 117-127.

931 Flanagan, L.B., Brooks, J.R., Varney, G.T., Berry, S.C., Ehleringer, J.R., 1996. Carbon isotope
 932 discrimination during photosynthesis and the isotope ratio of respired CO_2 in boreal forest
 933 ecosystems. *Global Biogeochemical Cycles* 10(4), 629-640.

934 Fletcher, B.J., Beerling, D.J., Chaloner, W., 2004. Stable carbon isotopes and the metabolism of
 935 the terrestrial Devonian organism *Spongiophyton*. *Geobiology* 2, 107-119.

936 Friedman, G.M., Sanders, J.E., 1982. Time-temperature-burial significance of Devonian
 937 anthracite implies former great (~6.5 km) depth of burial of Catskill Mountains, New York.
 938 Geology 10, 93-96.

939 Gastaldo, R.A., 2016. New paleontological insights into the Emsian–Eifelian Trout Valley
 940 Formation, Baxter State Park's Scientific Forest Management Area, Aroostock County,
 941 Maine. Palaios 31(7), 339-346.

942 Gensel, P.G., 1992. Phylogenetic relationships of the zosterophylls and lycopsids: evidence from
 943 morphology, paleoecology, and cladistics methods of inference. Ann. Missouri Bot. Gard.
 944 79(3), 450-473.

945 Gensel, P.G., 2017. Silurian origins of ferns and lycophytes - what we know, what we need to
 946 find out. Fern Gazette 20(6), 217-242.

947 Gensel, P.G., Albright, V.M., 2006. *Leclercqia complexa* from the Early Devonian (Emsian) of
 948 northern New Brunswick, Canada. Rev. Palaeobot. Palynol. 142, 103-121.

949 Gensel, P.G., Andrews, H.N., 1984. Plant Life in the Devonian. Praeger, New York, 380 pp.

950 Gensel, P.G., Berry, C.M., 2001. Early lycophyte evolution. Am. Fern J. 91(3), 74-98.

951 Gensel, P.G., Berry, C.M., 2016. Sporangial morphology of the Early Devonian zosterophyll
 952 *Sawdonia ornata* from the type locality (Gaspé). Internat. J. Plant Sci. 177(7), 618-632.

953 Gensel, P.G., Edwards, D. (Eds.), 2001. Plants Invade the Land: Evolutionary and Environmental
 954 Perspectives. Columbia Univ. Press, 304 pp.

955 Gensel, P.G., Kasper, A.E., Jr., 2005. A new species of the Devonian lycopod genus, *Leclercqia*,
 956 from Emsian of New Brunswick, Canada. Rev. Palaeobot. Palynol. 137, 105-123.

957 Gensel, P.G., Pigg, K.B., 2010. An arborescent lycopsid from the Lower Carboniferous Price
958 Formation, southwestern Virginia, USA and the problem of species delimitation. Intern. J.
959 Coal Geol. 83(2-3), 132-145.

960 Gensel, P.G., Andrews, H.N., Forbes, W.H., 1975. A new species of *Sawdonia* with notes on the
961 origin of microphylls and lateral sporangia. Bot. Gaz. 136(1), 50-62.

962 Gensel, P.G., Kotyk, M., Basinger, J.M., 2001. Morphology of above- and below-ground
963 structures in Early Devonian plants. In: Gensel, P.G., Edwards, D. (Eds.), Plants Invade the
964 Land: Evolutionary and Environmental Perspectives. Columbia Univ. Press, New York, pp.
965 83-102.

966 Gerrienne, P., 1995. Les fossils végétaux du Dévonien inférieur de Marchin (bord nor du
967 Synclinorium de Dinant, Belgique). III. *Psilophyton parvulum* nov. sp. Geobios 28(2), 131-
968 144.

969 Gerrienne, P., 1997. The fossil plants from the Lower Devonian of Marchin (northern margin of
970 Dinant Synclinorium, Belgium). V. *Psilophyton genseliae* sp. Nov., with hypotheses on the
971 origin of Trimerophytina. Rev. Palaeobot. Palynol. 98, 303-324.

972 Gerrienne, P., Meyer-Berthaud, B., Yang, N., Steemans, P., Li, C.S., 2014. *Planatophyton* gen.
973 nov., a Middle Devonian euphyllrophyte from Xinjiang, north-west China. Rev. Palaeobot.
974 Palynol. 208, 55-64.

975 Gibling, M.R., Davies, N.S., 2012. Palaeozoic landscapes shaped by plant evolution. Nature
976 Geoscience 5(2), 99-105.

977 Gibling, M.R., Davies, N.S., Falcon-Lang, H.J., Bashforth, A.R., DiMichele, W.A., Rygel, M.C., Ielpi,
978 A., 2014. Palaeozoic co-evolution of rivers and vegetation: a synthesis of current
979 knowledge. *Proc. Geol. Assoc.* 125(5-6), 524-533.

980 Gradstein, F.M., Ogg, J.G., Schmitz, M., Ogg, G. (Eds.), 2012. *The Geologic Time Scale 2012*.
981 Elsevier, Amsterdam, 2 vols.

982 Graham, L.E., Gray, J., 2001. The origin, morphology, and ecophysiology of early embryophytes:
983 neontological and paleontological perspectives. In: Gensel, P.G., Edwards, D. (Eds.), *Plants*
984 *Invade the Land: Evolutionary and Environmental Perspectives*, Columbia Univ. Press, New
985 York, pp. 140-159.

986 Graham, L.E., Cook, M.E., Hanson, D.T., Pigg, K.B., Graham, J.M., 2010. Structural, physiological,
987 and stable carbon isotopic evidence that the enigmatic Paleozoic fossil *Prototaxites* formed
988 from rolled liverwort mats. *Am. J. Bot.* 97(2), 268-275.

989 Granoff, J.A., Gensel, P.G., Andrews, H.N., 1976. A new species of *Pertica* from the Devonian of
990 Canada. *Palaeontographica (B)* 155, 119-128.

991 Greb, S.F., DiMichele, M.A., Gastaldo, R.A., 2006. Evolution and importance of wetlands in earth
992 history. In: Greb, S.F., DiMichele, M.A. (Eds.), *Wetlands Through Time*, *Geol. Soc. Am. Spec.*
993 *Publ.* 399, pp. 1-40.

994 Grierson, J.D., Banks, H.P., 1983. A new genus of lycopods from the Devonian of New York
995 State. *Bot. J. Linn. Soc.* 86, 81-101.

996 Grierson, J.D., Bonamo, P.M., 1979. *Leclercqia complexa*: Earliest ligulate lycopod (Middle
997 Devonian). *Am. J. Bot.* 66, 474-476.

- 998 Griffing, D.H., Bridge, J.S., Hotton, C.L., 2000. Coastal-fluvial palaeoenvironments and plant
999 palaeoecology of the Lower Devonian (Emsian), Gaspé Bay, Quebec, Canada. In: Friend,
1000 P.F., Williams, B.P.J. (Eds.), New Perspectives on the Old Red Sandstone. Geol. Soc. London
1001 Spec. Publ. 180, pp. 61-84.
- 1002 Guo, J., Tan, H., Ding, S., Wang, S., 2010. Study on continuous carbonization process of biomass.
1003 www.igu.org/html/wgc2009/papers/docs/wgcFinal00123.pdf.
- 1004 Gupta, N.S., Briggs, D.E., Collinson, M.E., Evershed, R.P., Michels, R., Pancost, R.D., 2007.
1005 Molecular preservation of plant and insect cuticles from the Oligocene Enspel Formation,
1006 Germany: evidence against derivation of aliphatic polymer from sediment. Org.
1007 Geochem. 38(3), 404-418.
- 1008 Hammond, S.E., Berry, C.M., 2005. A new species of *Tetraxylopteris* (Aneurophytales) from the
1009 Devonian of Venezuela. Bot. J. Linn. Soc. 148, 275-303.
- 1010 Hao, S.G., Xue, J.Z., 2013a. Earliest record of megaphylls and leafy structures, and their initial
1011 diversification. Chinese Science Bulletin 58(23), 2784-2793.
- 1012 Hao, S.G., Xue, J.Z., 2013b. The Early Devonian Posongchong Flora of Yunnan. Science Press,
1013 Beijing, 366 pp.
- 1014 Heroux, Y., Chagnon, A., Bertrand, R., 1979. Compilation and correlation of major thermal
1015 maturation indicators. Am. Assoc. Petrol. Geol. Bull. 63(12), 2128-2144.
- 1016 Hilton, J., Bateman, R.M., 2006. Pteridosperms are the backbone of seed-plant phylogeny. J.
1017 Torrey Bot. Soc. 133(1), 119-168.
- 1018 Hotton, C.L., Hueber, F.M., Griffing, D.H., Bridge, J.S., 2001. Early terrestrial plant environments:
1019 An example from the Emsian of Gaspé, Canada. In: Gensel, P.G., Edwards, D. (Eds.), Plants

1020 Invade the Land: Evolutionary and Environmental Perspectives, Columbia Univ. Press, New
1021 York, pp. 179-212.

1022 Hueber, F.M., 1971. *Sawdonia ornata*: A new name for *Psilophyton princeps* var. *ornatum*.
1023 Taxon 20, 641-642.

1024 Hueber, F.M., Grierson, J.D., 1961. On the occurrence of *Psilophyton princeps* in the early Upper
1025 Devonian of New York. Am. J. Bot. 48, 473-479.

1026 Jackson, P.C., Meinzer, F.C., Goldstein, G., Holbrook, N.M., Cavelier, J., Rada, F., 1993.
1027 Environmental and physiological influences on carbon isotope composition of gap and
1028 understory plants in a lowland tropical forest. In: Ehleringer, J.R., Hall, A.E., Farquhar, G.D.
1029 (Eds.), Stable Isotopes and Plant Carbon-Water Relations, Academic Press, pp. 131-140.

1030 Jennings, J., 1976. The morphology and relationships of *Rhodea*, *Telangium*, *Telangiopsis*, and
1031 *Heterangium*. Am. J. Bot. 63(8), 1119-1133.

1032 Kasper, A.E., Jr., Andrews, H.N., Jr., 1972. *Pertica*, a new genus of Devonian plants from
1033 northern Maine. Am. J. Bot. 59(9), 897-911.

1034 Kennedy, K., Gensel, P.G., Gibling, M.R., 2012. Paleoenvironmental inferences from the classic
1035 Early Devonian plant-bearing locality of the Campbellton Formation, New Brunswick,
1036 Canada. Palaios 27, 424-438.

1037 Kennedy, K.L., Gibling, M.R., Eble, C.F., Gastaldo, R.A., Gensel, P.G., Werner-Zwanziger, U.,
1038 Wilson, R.A., 2013. Lower Devonian coaly shales of northern New Brunswick, Canada: plant
1039 accumulations in the early stages of terrestrial colonization. J. Sed. Res. 83(12), 1202-1215.

1040 Kenrick, P., Crane, P.R., 1997. The origin and early evolution of plants on land. Nature 389, 33-
1041 39.

1042 Knaus, M.J., 1994. *Triphylopteris collombiana*: a clarification of the generic concepts based on
1043 rediscovered specimens from Kossberg bei Plauen, Germany, and a reassignment of the
1044 North American species of *Triphylopteris* to *Genselia* gen. nov. Int. J. Plant Sci. 155(1), 97-
1045 116.

1046 Knaus, M.J., 1995. The species of the Early Carboniferous fossil plant genus *Genselia*. Int. J.
1047 Plant Sci. 156(1), 61-92.

1048 Kohn, M.J., 2010. Carbon isotope compositions of terrestrial C3 plants as indicators of
1049 (paleo)ecology and (paleo)climate. Proc. Nat. Acad. Sci. USA 107(46), 19691-19695.

1050 Lawrence, D., 1986. Sedimentology of the Lower Devonian Battery Point Formation, eastern
1051 Gaspé Peninsula. Ph.D. thesis, University of Bristol, Bristol, U.K.

1052 Leclercq, S., Banks, H.P., 1962. *Pseudosporochnus nodosus* sp. nov., a Middle Devonian plant
1053 with cladoxylalean affinities. Palaeontographica (B) 110, 1-34.

1054 Le Hir, G., Donnadieu, Y., Godd  ris, Y., Meyer-Berthaud, B., Ramstein, G., Blakey, R.C., 2011.
1055 The climate change caused by the land plant invasion in the Devonian. Earth Planet. Sci.
1056 Lett. 310, 203-212.

1057 Li, C.S., Edwards, D., 1995. A re-investigation of Halle's *Drepanophycus spinaeformis* G  pp from
1058 the Lower Devonian of Yunnan Province, southern China. Bot. J. Linn. Soc. 118, 163-192.

1059 Li, C.S., Hueber F.M., Hutton C.L., 2000. A neotype for *Drepanophycus spinaeformis* Goppert
1060 1852. Can. J. Bot. 78, 889-902.

1061 Liu, W., Feng, X., Ning, Y., Zhang, Q., Cao, Y., An, Z., 2005. $\delta^{13}\text{C}$ variation of C3 and C4 plants
1062 across an Asian monsoon rainfall gradient in arid northwestern China. Glob. Change Biol.
1063 11, 1094-1100.

1064 Macfarlane, C., Adams, M.A., White, D.A., 2004. Productivity, carbon isotope discrimination and
1065 leaf traits of trees of *Eucalyptus globulus* Labill in relation to water availability. Plant Cell
1066 Environ. 27, 1515-1524.

1067 Malamud-Roam, F., Ingram, B.L., 2001. Carbon isotopic compositions of plants and sediments
1068 of tide marshes in the San Francisco estuary. J. Coast. Res. 17(1), 17-29.

1069 Marino, B.D., McElroy, M.B., 1991. Isotopic composition of atmospheric CO₂ inferred from in C₄
1070 plant cellulose. Nature, 127-131.

1071 Marshall, J.E.A., 1996. *Rhabdosporites langii*, *Geminospora lemurata* and *Contagisporites*
1072 *optivus*: An origin for heterospory within the progymnosperms. Rev. Palaeobot. Palynol. 93,
1073 159-189.

1074 Meyer-Berthaud, B., Scheckler, S.E., Wendt, J., 1999. *Archaeopteris* is the earliest known
1075 modern tree. Nature 398, 700-704.

1076 Meyer-Berthaud, B., Fairon-Demaret, M., Steemans, P., Talent, J., Gerrienne, P., 2003. The plant
1077 *Leclercqia* (Lycopsida) in Gondwana: implications for reconstructing Middle Devonian
1078 palaeogeography. Geol. Mag. 140, 119-130.

1079 Meyer-Berthaud, B., Soria, A., Decombeix, A.L., 2010. The land plant cover in the Devonian: A
1080 reassessment of the evolution of the tree habit. In: Vecoli, M., Clément, G., Meyer-
1081 Berthaud, B. (eds.), The Terrestrialization Process: Modelling Complex Interactions at the
1082 Biosphere-Geosphere Interface, Geological Society of London, Spec. Publ. 339, pp. 59-70.

1083 Milici, R.C., Swezey, C.S., 2006. Assessment of Appalachian basin oil and gas resources;
1084 Devonian Shale-Middle and Upper Paleozoic total petroleum system: U.S. Geological
1085 Survey Open-File Report 2006-1237, 70 pp. (<http://pubs.usgs.gov/of/2006/1237/>).

1086 Miller, J.M., Williams, R.J., Farquhar, G.D., 2001. Carbon isotope discrimination by a sequence
1087 of *Eucalyptus* species along a subcontinental rainfall gradient in Australia. *Funct. Ecol.* 15,
1088 222-232.

1089 Mintz, J.S., Driese, S.G., White, J.D., 2010. Environmental and ecological variability of Middle
1090 Devonian (Givetian) forests in Appalachian Basin paleosols, New York, United States.
1091 *Palaios* 25, 85-96.

1092 Mora, C.I., Driese, S.G., Colarusso, L.A., 1996. Middle to late Paleozoic atmospheric CO₂ levels
1093 from soil carbonate and organic matter. *Science* 271(5252), 1105-1107.

1094 Nilsen, E., Sharifi, M., 1997. Carbon isotopic composition of legumes with photosynthetic stems
1095 from mediterranean and desert habitats. *Am. J. Bot.* 84(12), 1707-1707.

1096 Obermajer, M., Fowler, M.G., Goodarzi, F., Snowdon, L.R., 1997. Organic petrology and organic
1097 geochemistry of Devonian black shales in southwestern Ontario, Canada. *Org. Geochem.*
1098 26, 229-246.

1099 Obukhovskaya, T., 2000. Miospores of the Givetian-Frasnian boundary deposits in Belarus. *Acta*
1100 *Palaeobotanica* 40(1), 17-23.

1101 O'Leary, M.H., 1988. Carbon isotopes in photosynthesis. *BioScience* 38(5), 328-336.

1102 Pate, J., Arthur, D., 1998. $\delta^{13}\text{C}$ analysis of phloem sap carbon: Novel means of evaluating
1103 seasonal water stress and interpreting carbon isotope signatures of foliage and trunk wood
1104 of *Eucalyptus globulus*. *Oecologia* 117, 301-311.

1105 Playford, G., Dino, R., 2005. Carboniferous and Permian palynostratigraphy. In: Koutsoukos,
1106 E.A.M. (Ed.), *Applied Stratigraphy*. Springer, Amsterdam, pp. 101-121.

1107 Poss, J.A., Grattan, S.R., Suarez, D.L., Grieve, C.M., 2000. Stable carbon isotope discrimination:
 1108 an indicator of cumulative salinity and boron stress in *Eucalyptus camaldulensis*. Tree Phys.
 1109 20, 1121-1127.

1110 Pratt, B.R., van Heerde, J., 2017. An arborescent lycopsid stem fragment from the Palliser
 1111 Formation (Famennian) carbonate platform, southwestern Alberta, Canada, and its
 1112 paleogeographic and paleoclimatic significance. Can. J. Earth Sci. 54(2), 141-145.

1113 Prestianni, C., Meyer-Berthaud, B., Blanchard, R., Rücklin, M., Clément, G., Gerrienne, P., 2012.
 1114 The Middle Devonian plant assemblage from Dechra Aït Abdallah (Central Morocco)
 1115 revisited. Rev. Palaeobot. Palynol. 179, 44-55.

1116 Raich, J.W., Tufekciogul, A., 2000. Vegetation and soil respiration: correlations and controls.
 1117 Biogeochemistry 48(1), 71-90.

1118 Raven, P.H., Evert, R.F., Eichhorn, S.E. (Eds.), 2004. Biology of Plants, 7th ed. W.H. Freeman, New
 1119 York, 686 pp.

1120 Raymond, A., Metz, C., 1995. Laurussian land-plant diversity during the Silurian and Devonian:
 1121 mass extinction, sampling bias, or both? Paleobiology 21, 74-91.

1122 Rayner, R.J., 1983. New observations on *Sawdonia ornata* from Scotland. Trans. Roy. Soc.
 1123 Edinburgh (Earth Sciences) 74, 79-93.

1124 Read, C.B., 1955. Floras of the Pocono Formation and Price Sandstone in parts of Pennsylvania,
 1125 Maryland, West Virginia, and Virginia. U.S. Geol. Surv. Prof. Paper 263, 30 pp.

1126 Repetski, J.E., Ryder, R.T., Weary, D.J., Harris A.G., Trippi, M.H., 2008. Thermal maturity
 1127 patterns (CAI and %R₀) in Upper Ordovician and Devonian rocks of the Appalachian Basin:

1128 a major revision of USGS Map I-917-E using new subsurface collections. U.S. Geol. Survey,
1129 Sci. Inv. Map 3006.

1130 Retallack, G.J., Huang, C., 2011. Ecology and evolution of Devonian trees in New York, USA.
1131 *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 299, 110-128.

1132 Richardson, J.B., McGregor, D.C., 1986. Silurian and Devonian spore zones of the Old Red
1133 Sandstone continent and adjacent regions. *Geol. Surv. Can. Bull.* 384, 1-79.

1134 Roden, J.S., Bowling, D.R., McDowell, N.G., Bond, B.J., Ehleringer, J.R., 2005. Carbon and oxygen
1135 isotope ratios of tree ring cellulose along a precipitation transect in Oregon, United States.
1136 *J. Geophys. Res.* 110, G02003, 11 pp., doi: 10.1029/2005JG000033.

1137 Rowan, E.L., 2006. Burial and thermal history of the Central Appalachian Basin, based on three
1138 2-D models of Ohio, Pennsylvania, and West Virginia. U.S. Geol. Surv., Open-File Report
1139 2006-1019.

1140 Rygel, M.C., Calder, J.H., Gibling, M.R., Gingras, M.K., Melrose, C.S.A., 2006. Tournaisian
1141 forested wetlands in the Horton Group of Atlantic Canada. In: Greb, S.F., DiMichele, M.A.
1142 (Eds.), *Wetlands Through Time*. *Geol. Soc. Am. Spec. Publ.* 399, pp. 103-126.

1143 Scheckler, S.E., 1978. Ontogeny of progymnosperms II. Shoots of Upper Devonian
1144 *Archaeopteridales*. *Can. J. Bot.* 56, 3136-3170.

1145 Scheckler, S.E., 1986a. Geology, floristics and paleoecology of Late Devonian coal swamps from
1146 Appalachian Laurentia (U.S.A.). *Ann. Soc. Géol. Belg.* 109, 209-222.

1147 Scheckler, S.E., 1986b. Floras of the Devonian–Mississippian transition. In: Gastaldo, R.A. (Org.)
1148 & Broadhead, T.W. (Ed.), *Land Plants: Notes for a Short Course*. Univ. Tenn. Dept. Geol. Sci.,
1149 Knoxville, Tenn., pp. 81-96.

1150 Scheckler, S.E., 2001. Afforestation—The first forests. In: Briggs, D.E.G., Crowther, P. (Eds.),
1151 Palaeobiology II. Blackwell, Oxford, pp. 67–71.

1152 Scheckler, S.E., Banks, H.P., 1971. Anatomy and relationships of some Devonian
1153 progymnosperms from New York. *Am. J. Bot.* 58(8), 737-751.

1154 Schimmelmann, A., Qi, H., Coplen, T.B., Brand, W.A., Fong, J., Meier-Augenstein, W., Kemp,
1155 H.F., Toman, B., Ackermann, A., Assonov, S., Aerts-Bijma, A.T., 2016. Organic reference
1156 materials for hydrogen, carbon, and nitrogen stable isotope-ratio measurements: caffeines,
1157 n-alkanes, fatty acid methyl esters, glycines, L-valines, polyethylenes, and oils. *Anal. Chem.*
1158 88(8), 4294-4302.

1159 Schleser, G.H., Jayasekera, R., 1985. $\delta^{13}\text{C}$ -variations of leaves in forests as an indication of
1160 reassimilated CO_2 from the soil. *Oecologia* 65(4), 536-542.

1161 Silvestro, D., Cascales-Miñana, B., Bacon, C.D., Antonelli, A., 2015. Revisiting the origin and
1162 diversification of vascular plants through a comprehensive Bayesian analysis of the fossil
1163 record. *New Phytol.* 207(2), 425-436.

1164 Skog, J.E., Gensel, P.G., 1980. A fertile species of *Triphyllopteris* from the Early Carboniferous
1165 (Mississippian) of Southwestern Virginia. *Am. J. Bot.* 67(4), 440-451.

1166 Sperry, J.S., 2003. Evolution of water transport and xylem structure. *Internat. J. Plant Sci.*
1167 164(S3), S115-S127.

1168 Spicer, R.A., 1989. Physiological characteristics of land plants in relation to environment
1169 through time. *Trans. Roy. Soc. Edinburgh (Earth Sci.)* 80, 321-9.

1170 Stein, W., 1993. Modeling the evolution of stelar architecture in vascular plants. *Int. J. Plant Sci.*
1171 154(2), 229-263.

1172 Stein, W.E., Mannolini, F., Hernick, L.V., Landing, E., Berry, C.M., 2007. Giant cladoxylopsid trees
1173 resolve the enigma of the Earth's earliest forest stumps at Gilboa. *Nature* 446(7138), 904-
1174 907.

1175 Stein, W.E., Berry, C.M., VanAller Hernick, L., Mannolini, F., 2012. Surprisingly complex
1176 community discovered in the mid-Devonian fossil forest at Gilboa. *Nature* 483, 78-81.

1177 Stewart, W.N., Rothwell, G.W., 1993. *Paleobotany and the Evolution of Plants*, 2nd ed.
1178 Cambridge Univ. Press, Cambridge, 521 pp.

1179 Streel, M., Caputo, M.V., Loboziak, S., Melo, J.H.G., 2000. Late Frasnian-Famennian climates
1180 based on palynomorphs analyses and the question of the Late Devonian glaciations. *Earth-*
1181 *Sci. Rev.* 52, 121-173.

1182 Stubblefield, S., Banks H.P., 1978. The cuticle of *Drephanophycus spinaeformis*, a long-ranging
1183 Devonian Lycopod from New York and Eastern Canada. *Am. J. Bot.* 65(1), 110-118.

1184 Taylor, T.N., Taylor, E.L., Krings, M. (Eds.), 2009. *Paleobotany: The Biology and Evolution of*
1185 *Fossil Plants*, 2nd ed. Academic Press, London, 1230 pp.

1186 Trant, C.A., Gensel, P.G., 1985. Branching in *Psilophyton*: a new species from the Lower
1187 Devonian of New Brunswick, Canada. *Am. J. Bot.* 72(8), 1256-1273.

1188 Turnau, E., 2014. Floral change during the Taghanic Crisis: spore data from the Middle Devonian
1189 of northern and south-eastern Poland. *Rev. Palaeobot. Palynol.* 200, 108-121.

1190 Utting, J., Hamblin, A.P., 1991. Thermal maturity of the Lower Carboniferous Horton Group,
1191 Nova Scotia. *Int. J. Coal Geol.* 19, 439-456.

1192 van der Voo, R., 1988. Paleozoic paleogeography of North America, Gondwana, and intervening
1193 displaced terranes: Comparisons of paleomagnetism with paleoclimatology and
1194 biogeographical patterns. Geol. Soc. Am. Bull. 100, 311-324.

1195 van der Voo, R., 1993. Paleomagnetism of the Atlantic, Tethys, and Iapetus Oceans. Cambridge
1196 Univ. Press, Cambridge, 411 pp.

1197 Wellman, C.H., Osterloff, P.L., Mohiuddin, U., 2003. Fragments of the earliest land plants.
1198 Nature 425(6955), 282-285.

1199 Wilson, J.P., 2013. Modeling 400 million years of plant hydraulics. Paleontol. Soc. Pap. 19, 175–
1200 194.

1201 Wilson, J.P., 2016. Hydraulics of *Psilophyton* and evolutionary trends in plant water transport
1202 after terrestrialization. Rev. Palaeobot. Palynol. 227, 65-76.

1203 Wilson, J.P., Fischer, W.W., 2011. Hydraulics of *Asteroxylon mackiei*, an early Devonian vascular
1204 plant, and the early evolution of water transport tissue in terrestrial plants. Geobiology 9,
1205 121–130.

1206 Wilson, J.P., Knoll, A.H., 2010. A physiologically explicit morphospace for tracheid-based water
1207 transport in modern and extinct seed plants. Paleobiology 36, 335–355.

1208 Winter, K., Holtum, J.A.M., 2005. The effects of salinity, crassulacean acid metabolism and plant
1209 age on the carbon isotope composition of *Mesembryanthemum crystallinum* L., a
1210 halophytic C3-CAM species. Planta 222, 201-209.

1211 Xu, H.H., Berry, C.M., 2008. The Middle Devonian lycopsid *Haskinsia* Grierson et Banks from the
1212 Ruppert Coast, Marie Byrd Land, West Antarctica. Rev. Palaeobot. Palynol. 150, 1-4.

1213 Xu, H.H., Wang, Y., 2008. The palaeogeographical significance of specimens attributed to
 1214 *Protolpidodendron scharyanum* Krejčí (Lycopsida) from the Middle Devonian of North
 1215 Xinjiang, China. Geological Magazine 145(2), 295-299.

1216 Xu, H.H., Wang, Y., Berry, C.M., Cai, C.Y., 2008. Two species of *Haskinsia* Grierson & Banks
 1217 (Lycopsida) from the Middle Devonian of Xinjiang, China, and consideration of their
 1218 palaeogeographical significance. Bot. J. Linn. Soc. 157, 633-644.

1219 Xu, H.H., Berry, C.M., Wang, Y., Marshall, J.E.A., 2011. A new species of *Leclercqia* Banks,
 1220 Bonamo et Grierson (Lycopsida) from the Middle Devonian of North Xinjiang, China, with a
 1221 possible climbing habit. Int. J. Plant Sci. 172, 836-846.

1222 Xu, H.H., Feng, J., Jiang, Q., Wang, Y., 2013. Report of *Drepanophycus* Göppert (Lycopsida) from
 1223 the Middle Devonian of Xinjiang, China. J. Syst. Evol. 51(6), 765-772.

1224 Xu, H.H., Jiang, Q., Zhang, X.L., Wang, Y., Feng, J., 2015. On the Mid Devonian Hujiersite flora
 1225 from West Junggar, Xinjiang, China, its characteristics, age, palaeoenvironment and
 1226 palaeophytogeographical significances. Acta Palaeontol. Sin. 54, 230-239.

1227 Xu, H.H., Berry, C.M., Stein, W., Wang, Y., Tang, P., Fu, Q., 2017. Unique growth strategy in the
 1228 Earth's first trees revealed in silicified fossil trunks from China. Proc. Nat. Acad. Sci. (U.S.A.)
 1229 114, 12009-12014.

1230 Xue, J., 2012. Lochkovian plants from the Xitun Formation of Yunnan, China, and their
 1231 palaeophytogeographical significance. Geol. Mag. 149(2), 333-344.

1232 Xue, J., Deng, Z., Huang, P., Huang, K., Benton, M.J., Cui, Y., Wang, D., Liu, J., Shen, B., Basinger,
 1233 J.F., Hao, S., 2016. Belowground rhizomes in paleosols: The hidden half of an Early
 1234 Devonian vascular plant. Proc. Nat. Acad. Sci. (U.S.A.) 113(34), 9451-9456.

1235 Xue, J., Huang, P., Wang, D., Xiong, C., Liu, L., Basinger, J.F., 2018. Silurian-Devonian terrestrial
1236 revolution in South China: Taxonomy, diversity, and character evolution of vascular plants
1237 in a palaeogeographically isolated, low-latitude region. *Earth-Sci. Rev.* 180, 92-125.

1238 Yang, N., Edwards, D., Ablaev, A., Li, C.S., 2008. The discovery of fertile *Haskinsia colophylla*
1239 (*Haskinsiaceae*) in Far East Russia. *Taxon* 57(2), 588-593.

1240 Zaiontz, C., 2014. Real statistics using Excel. Online. Tersedia di: [http://www. realstatistics.](http://www.realstatistics.com/tests-normality-and-symmetry/statistical-tests-normalitysymmetry/kolmogorov-smirnov-test/)
1241 [com/tests-normality-and-symmetry/statistical-tests-normalitysymmetry/kolmogorov-](http://www.realstatistics.com/tests-normality-and-symmetry/statistical-tests-normalitysymmetry/kolmogorov-smirnov-test/)
1242 [smirnov-test/](http://www.realstatistics.com/tests-normality-and-symmetry/statistical-tests-normalitysymmetry/kolmogorov-smirnov-test/)[diakses pada 03 Juni 2014] Lampiran, 1.

1243 Zheng, D., Xu, H., Wang, J., Feng, C., Zhang, H., Chang, S.C., 2016. Geochronologic age
1244 constraints on the Middle Devonian Hujiersite flora of Xinjiang, NW China. *Palaeogeogr.*
1245 *Palaeoclimatol. Palaeoecol.* 463, 230-237.

1246 Ziegler, W., Sandberg, C.A., 1990. The Late Devonian Standard Conodont Zonation. *Cour.*
1247 *Forschungsinst. Senckenberg* 121, 1-115.

1249 **Tables**

1250

1251 Table 1. Summary of carbon isotope analyses by taxon and plant part

1252

1253 Table 2. Mean differences in $\Delta^{13}\text{C}_p$ by taxon

1254

1255 Table 3. Differences in $\delta^{13}\text{C}_p$ between anatomical parts of individual specimens

1256

1257 Table S1. Stable carbon isotope compositions of Devonian-Carboniferous fossil plant specimens

1258

Table 1. Summary of carbon isotope analyses by taxon and plant part

	Number of specimens	Number of carbon isotope analyses					Mean $\delta^{13}\text{C}$				
		Stem	Leaf	Spine	Indet.	Total	Stem	Leaf	Spine	Indet.	Avg.
<i>Archaeopteris</i>	51	36	17		27	80	-25.53	-25.32		-27.16	-26.04
<i>Drepanophycus</i>	21	23		7	10	40	-26.08		-25.52	-25.90	-25.94
<i>Genselia</i>	9	2	1		7	10	-22.66	-23.08		-23.72	-23.44
<i>Haskinsia</i>	4	2			2	4	-22.27			-22.31	-22.29
<i>Leclercqia</i>	13	16		3	5	24	-22.96		-23.91	-22.71	-23.02
<i>Pertica</i>	11	12			8	20	-25.55			-25.26	-25.43
<i>Wattieza</i>	8	6			5	11	-26.21			-26.74	-26.45
<i>Psilophyton</i>	17	13			12	25	-24.46			-25.27	-24.85
<i>Rhacophyton</i>	18	19	4		10	33	-24.51	-25.28		-25.82	-25.00
<i>Rhodeopteridium</i>	3	2	1			3	-23.71	-22.82			-23.41
<i>Sawdonia</i>	14	18		4	6	28	-27.46		-27.08	-25.85	-27.07
<i>Tetraxylopteris</i>	21	19	2		10	31	-25.56	-26.09		-27.01	-26.06
Total / Average	190	168	25	14	102	309	-25.30	-25.18	-25.62	-25.87	-25.50

1259 Notes: Indet. = indeterminate. Avg. = average (of stem + leaf + spine + indet. mean values).

1260

Table 2. Mean differences in $\Delta^{13}\text{C}_p$ by taxon

	Number of analyses	mean (‰)	st. dev. (‰)	s.e. (‰)	Student's t	p(α)
<i>Archaeopteris</i>	80	-0.38	1.18	0.13	3.05	0.002
<i>Drepanophycus</i>	40	-0.26	1.08	0.17	1.32	0.189
<i>Genselia</i>	10	0.37	1.12	0.36	0.69	0.492
<i>Haskinsia</i>	4	2.41	1.48	0.74	3.31	0.001
<i>Leclercqia</i>	24	1.81	1.02	0.21	6.57	0.001
<i>Pertica</i>	20	0.40	0.56	0.12	1.06	0.292
<i>Wattieza</i>	11	-0.51	0.49	0.15	1.28	0.201
<i>Psilophyton</i>	25	0.94	1.55	0.31	3.22	0.001
<i>Rhacophyton</i>	33	0.23	1.32	0.23	0.65	0.517
<i>Rhodeopteridium</i>	3	0.41	0.89	0.51	0.42	0.677
<i>Sawdonia</i>	28	-1.22	1.81	0.34	5.02	0.001
<i>Tetraxylopteris</i>	31	-0.05	1.35	0.24	0.37	0.714

Notes: bold type denotes significant values ($= p(\alpha) < 0.05$); st. dev. = standard deviation;

s.e. = standard error of the mean. Statistical software from [Zaiontz \(2014\)](#).

1261

1262

Table 3. Differences in $\delta^{13}\text{C}_p$ between anatomical parts of individual specimens

	No. of pairs	Stem $\delta^{13}\text{C}$ (‰)	Leaf/spine $\delta^{13}\text{C}$ (‰)	Difference $\Delta^{13}\text{C}$ (‰)	s.e. $\delta^{13}\text{C}$ (‰)	Student's t	$p(\alpha)$
<i>Archaeopteris</i>	13	-0.08	0.33	0.42	0.20	1.54	0.137
<i>Drepanophycus</i>	6	-0.81	-0.12	0.69	0.30	1.15	0.277
<i>Leclercqia</i>	3	1.75	1.96	0.21	0.22	0.45	0.676
<i>Rhacophyton</i>	4	0.51	0.05	-0.46	0.23	1.54	0.173
<i>Sawdonia</i>	4	-2.77	-1.22	1.56	0.21	2.85	0.029
<i>Tetraxylopteris</i>	2	0.10	-0.06	-0.16	0.13	1.25	0.337
Total	32	-0.30	0.15	0.45	0.14	2.64	0.010

Notes: bold type denotes significant values ($=p(\alpha) < 0.05$); s.e. = standard error of the mean.

Figure captions

Figure 1. Devonian plant taxa of the present study: (A) phylogenetic relationships, and (B) morphological reconstructions. Reconstructions show full shoot system of plant except for *Tetraxylopteris* (lateral branch); open scale bars equal 10 cm and hachured scale bars equal 1 m. Sources: (A) Kenrick and Crane (1997), Hilton and Bateman (2006), and Taylor et al. (2009); (B) *Sawdonia* (<http://www.ucmp.berkeley.edu/IB181/VPL/Lyco/Lyco1.html>), *Drepanophycus* (Stewart and Rothwell, 1993), *Leclercqia* (Bonamo et al., 1988), *Haskinsia* (<http://dinoera.com/tags/haskinsia>), *Psilophyton* (<https://openclipart.org/detail/231687/psilophyton-princeps>), *Pertica* (Kasper and Andrews, 1972), *Wattieza* (Stein et al., 2007) [note: reconstructions of European *Pseudosporochnus* are slightly different, e.g., Berry and Fairon-Demaret, 2002], *Rhacophyton* (Cornet et al., 1977), *Rhodeopteridium* (Jennings, 1976), *Tetraxylopteris* (Bonamo and Banks, 1967), *Archaeopteris* (www.devoniantimes.org), and *Genselia* (Knaus, 1995).

Figure 2. Stratigraphic ages of fossil plant locales and ranges of fossil plant taxa used in this study. Marine conodont and terrestrial palynomorph zonation are from Richardson and McGregor (1986), Ziegler and Sandberg (1990), Streel et al. (2000), and Playford and Dino (2005); the timescale is from Gradstein et al. (2012). Locales: Gaspé Peninsula, Quebec, 1 = Haldimand Head, 2 = Aiguillon, Cap-aux-Os, and L'Anse-a-brilliant, and 3 = Seal Rock; New Brunswick, 1 = south shore Chaleur Bay (localities A, B, F), Dalhousie, Pin Sec Point, and

Peuplier Point, and 2 = Atholville and Restigouche; Hujiersite Fm (Xinjiang, China), 1 = Gannaren, 2 = Hwy 217, 3 = 251 Hill, and 4 = Hujiersite. Specimens of the Oneonta, Sonyea, Escuminac, and Pocono formations (*italicized*) are from multiple locales, but all fall within the narrow age ranges shown for each unit. For plant taxa, solid bars indicate peak abundance, open bars the full established biorange, and dashed bars uncertain extensions of the biorange.

Figure 3. Geographic distribution of fossil plant locales, shown on Late Devonian paleogeographic maps of (A) the world and (B) the Appalachian Basin. (A) Inset rectangle represents area of map B; abbreviations: A = Alberta, G = Germany, O = Ontario, V = Venezuela, X = Xinjiang, and Y = Yunnan; base map courtesy of Ron Blakey (Deep-Time Maps). Note that Laurentia (Paleozoic North America) was rotated ~30° clockwise relative to its present-day orientation ([van der Voo, 1988, 1993](#)). (B) Main fossil plant locales in Appalachian Basin; site names are positioned either directly above or below site locales (black dots); county names are given where multiple locales are within a small area. Dark brown field shows approximate area of basinward progradation of shoreline from the Early to the Late Devonian. N = present-day north; N_D = Devonian north.

Figure 4. Representative study specimens: (A) *Psilophyton forbesii* stems and leaves, Emsian, Gaspé Peninsula, Québec; (B) *Archaeopteris* sp. stems and leaves, Frasnian, West Virginia; (C) tracheids of *Psilophyton charientos*, Emsian, New Brunswick; (D) cuticle of *Sawdonia* sp., Emsian, Gaspé Peninsula, Québec. All specimens were inspected using a binocular scope to

verify the state of fossil preservation, and a subset was examined using scanning electron microscopy.

Figure 5. Model of $\Delta\delta^{13}\text{C}_{\text{p-X}}$ calculations. Relative to the LOWESS curve generated from the full plant $\delta^{13}\text{C}$ dataset, some fossil plant taxa exhibit systematically higher or lower $\delta^{13}\text{C}$ values. In this example, Taxon A exhibits higher-than-average $\delta^{13}\text{C}$ (positive $\Delta\delta^{13}\text{C}_{\text{p-X}}$), Taxon B exhibits average $\delta^{13}\text{C}$ (near-zero $\Delta\delta^{13}\text{C}_{\text{p-X}}$), and Taxon C exhibits lower-than-average $\delta^{13}\text{C}$ (negative $\Delta\delta^{13}\text{C}_{\text{p-X}}$). The complete LOWESS curve for the $\delta^{13}\text{C}_{\text{p}}$ dataset will be published in a companion study ([Algeo et al., in preparation](#)).

Figure 6. Deviations of $\delta^{13}\text{C}$ of Devonian fossil plant specimens from the LOWESS reference curve ($\Delta\delta^{13}\text{C}_{\text{p-X}}$) plotted as a function of specimen age. For each genus, the regression of $\Delta\delta^{13}\text{C}_{\text{p-X}}$ against time is shown as a thin line; these regression lines are shown not to suggest that the $\Delta\delta^{13}\text{C}_{\text{p-X}}$ of a given taxon changes significantly with time (for most taxa, there is not enough data to generate a statistically significant time-dependent trend) but, rather, to show which taxa yield $\Delta\delta^{13}\text{C}_{\text{p-X}}$ values that are systematically higher or lower than the age-specific mean $\Delta\delta^{13}\text{C}_{\text{p-X}}$ values (0 ‰). Note that some taxa are systematically ^{13}C -enriched and others are systematically ^{13}C -depleted.

Figure 7. Deviations of $\delta^{13}\text{C}$ of Devonian fossil plants from the LOWESS reference curve ($\Delta\delta^{13}\text{C}_{\text{p-X}}$) by taxon. Open circles represent individual C-isotope analyses; red diamond, open rectangle, and horizontal bar represent the mean, standard error of the mean, and

standard deviation range, respectively, for each taxon. The dashed vertical line represents no offset from the LOWESS trend, and the yellow field represents +/-1 standard deviation for the dataset as a whole. Note that some taxa are systematically ^{13}C -enriched and others are systematically ^{13}C -depleted.

Figure 8. Reconstructions of the ecological distribution of the 10 Devonian vascular land plant taxa of the present study: (A) Early Devonian (Emsian), (B) Middle Devonian (Givetian), and (C) Late Devonian (late Frasnian-early Famennian). Note that actual Devonian landscapes included other vascular and non-vascular plant taxa not shown here; these reconstructions are intended to show only the relative habitat preferences of the 10 Devonian taxa of the present study. Note further that the proposed paleo-environmental distributions of these taxa shown here are consistent with, but not proven by, the available isotopic data, and that alternative models are possible. Further study of Devonian fossil plant taxa will be needed to accurately constrain their actual habitat preferences.






Figure 1







Figure 3





Figure 4




Figure 5




Figure 6




Figure 7

